arXiv:1610.05294v1 [math.DS] 17 Oct 2016

SIMPLE LYAPUNOV SPECTRUM FOR CERTAIN LINEAR COCYCLES OVER PARTIALLY HYPERBOLIC MAPS

MAURICIO POLETTI AND MARCELO VIANA

Abstract. Criteria for the simplicity of the Lyapunov spectra of linear cocycles have been found by Furstenberg, Guivarc'h-Raugi, Gol'dsheid-Margulis and, more recently, Bonatti-Viana and Avila-Viana. In all the cases, the authors consider cocycles over hyperbolic systems, such as shift maps or Axiom A diffeomorphisms.

In this paper we propose to extend such criteria to situations where the base map is just partially hyperbolic. This raises several new issues concerning, among others, the recurrence of the holonomy maps and the (lack of) continuity of the Rokhlin disintegrations of u-states.

Our main results are stated for certain partially hyperbolic skew-products whose iterates have bounded derivatives along center leaves. They allow us, in particular, to exhibit non-trivial examples of stable simplicity in the partially hyperbolic setting.

1. Introduction

The theory of linear cocycles is a classical and rather developed field of Dynamical Systems and Ergodic Theory, whose origins go back to the works of Furstenberg, Kesten [13], Furstenberg [11] and Oseledets [17]. The simplest examples are the derivative transformations of smooth dynamical systems but the notion of linear cocycle is a lot more general and flexible, and arises naturally in many other situations, e.g., the spectral theory of Schrödinger operators.

Among the outstanding issues is the problem of simplicity: *when is it the case that the dimension of all Oseledets subspaces is equal to 1?* This was first studied by Furstenberg [11], Guivarc'h-Raugi [15] and Gol'dsheid-Margulis [14], who obtained explicit simplicity criteria for random i.i.d. products of matrices. Bonatti-Viana [7] and Avila-Viana [3] have much extended the theory, to include a much broader class of (Hölder continuous) cocycles over hyperbolic maps.

Our purpose in this paper is to initiate the study of simplicity in the context of linear cocycles over partially hyperbolic maps. The study of partially hyperbolic systems was introduced in the works of Brin-Pesin [9] and Hirsch-Pugh-Shub [16] and has been at the heart of much recent progress in this area. While sharing many of the important features of uniformly hyperbolic systems, partially hyperbolic maps are a lot more flexible and encompass many new interesting phenomena.

As we are going to see, the study of linear cocycles over partially hyperbolic maps introduces a host of new issues. We will deal with these issues in the context when

Date: October 18, 2016.

2010 *Mathematics Subject Classification.* 37H15, 37D25, 37D30.

Key words and phrases. Lyapunov exponent, linear cocycle, partial hyperbolicity.

M. P. was supported through doctoral and post-doctoral scholarships from CNPq and CAPES. M. V. was partially supported by CNPq, FAPERJ and PRONEX.

the partially hyperbolic map is a partially hyperbolic skew-product $\hat{f} : \hat{\Sigma} \times K \to \hat{\Sigma} \times K$ (this and other notions mentioned in what follows are defined precisely in Section 2).

The case when the map is non-uniformly hyperbolic, that is, when all center Lyapunov exponents are non-zero, is better understood. Indeed, the case of 2-dimensional cocycles is covered by Viana [19] and for general $d \geq 2$ it should be possible to combine a symbolic description as in Sarig [18] with the main result of Avila, Viana [3], that deals with cocycles over countable shifts.

For this reason, we focus on the opposite case: namely, we take the skew-product to be mostly neutral along the center direction, meaning that its iterates have bounded derivatives along the verticals $\{\hat{x}\} \times K$.

Concerning the linear cocycle, we take it to admit strong-stable and strong-unstable holonomies, in the sense of Bonatti-Gomez-Mont-Viana [6] and Avila-Santamaria-Viana [2]. The simplicity conditions in our main result, that we are going to state next, may be viewed as extensions of the pinching and twisting conditions in Bonatti-Viana [7] and Avila-Viana [3] to the present partially hyperbolic setting.

Firstly, we call the linear cocycle *uniformly pinching* if there exists some fixed (or periodic) vertical leaf $\ell = \{\hat{p}\} \times K$ such that the restriction to ℓ of every exterior power of the cocycle admits an invariant dominated decomposition

$$\ell \times \Lambda^k(\mathbb{C}^d) = E^{k,1} \oplus \cdots \oplus E^{k,\dim \Lambda^k(\mathbb{C}^d)}$$

into 1-dimensional subbundles. In particular, this decomposition is continuous and the Lyapunov exponents along the factor subbundles E^j are all distinct.

Secondly, we say that the linear cocycle is *uniformly twisting* if, for any *su*-path γ connecting points $(\hat{p}, t) \in \ell$ and $(\hat{p}, s) \in \ell$, the push-forward of the decomposition

$$E^{1,1}(t) \oplus \cdots \oplus E^{1,d}(t)$$

at $(\hat{p}, t)$ under the concatenation of strong-stable and strong-unstable holonomies over γ is in general position with respect to the decomposition

$$E^{1,1}(s) \oplus \cdots \oplus E^{1,d}(s)$$

at $(\hat{p}, s)$. By the latter, we mean that the image of any sum of k subspaces $E^{1,i}(t)$ is transverse to the sum of any $d - k$ subspaces $E^{1,j}(s)$, for any $1 \leq k \leq d - 1$.

We say that the linear cocycle is *uniformly simple* if it is both uniformly pinching and uniformly twisting.

Theorem A. Every uniformly simple linear cocycle over a partially hyperbolic skew-product with mostly neutral center direction has simple Lyapunov spectrum relative to any invariant probability measure with local partial product structure.

The notion of local partial product structure will be recalled in Section 2, where we will also give a more precise version of the theorem. Indeed, as we will see, the conclusion holds under weaker (non-uniform) versions of the pinching and twisting conditions. Moreover, already in the form given in Theorem A, our simplicity criterion holds for a subset of Hölder continuous cocycles with non-empty interior. We will also discuss the issue of density of this subset in Section 2.

Acknowledgements. We are grateful to L. Backes, F. Lenarduzzi, E. Pujals and J. Yang for numerous discussions.

2. Definitions and statements

Now let us give the precise definitions of the notions involved, as well as a refined version of Theorem A.

2.1. **Linear cocycles and Lyapunov exponents.** The *linear cocycle* defined by a measurable matrix-valued function $\hat{A} : \hat{M} \to GL(d, \mathbb{C})$ over an invertible measurable map $\hat{f} : \hat{M} \to \hat{M}$ is the (invertible) map $\hat{F}_A : \hat{M} \times \mathbb{C}^d \to \hat{M} \times \mathbb{C}^d$ given by

$$\hat{F}_A(\hat{x}, v) = \big(\hat{f}(\hat{x}), \hat{A}(\hat{x})v\big).$$

Its iterates are given by $\hat{F}_A^n(\hat{x}, v) = \big(\hat{f}^n(\hat{x}), \hat{A}^n(\hat{x})v\big)$ where

$$\hat{A}^n(\hat{x}) = \begin{cases} \hat{A}(\hat{f}^{n-1}(x)) \dots \hat{A}(\hat{f}(x))\hat{A}(\hat{x}) & \text{if } n > 0 \\ \mathrm{id} & \text{if } n = 0 \\ \hat{A}(\hat{f}^n(\hat{x}))^{-1} \dots \hat{A}(\hat{f}^{-1}(\hat{x}))^{-1} & \text{if } n < 0. \end{cases}$$

Let $\hat{\mu}$ be an $\hat{f}$-invariant probability measure on $\hat{M}$ such that $\log \|\hat{A}^{\pm 1}\|$ are integrable. By Oseledets [17], at $\hat{\mu}$-almost every point $\hat{x} \in \hat{M}$ there exist real numbers $\lambda_1(\hat{x}) > \cdots > \lambda_k(\hat{x})$ and a decomposition $\mathbb{C}^d = E^1_{\hat{x}} \oplus \cdots \oplus E^k_{\hat{x}}$ into vector subspaces such that

$$\hat{A}(\hat{x})E^i_{\hat{x}} = E^i_{\hat{f}(\hat{x})} \text{ and } \lambda_i(\hat{x}) = \lim_{|n| \to \infty} \frac{1}{n} \log \|\hat{A}^n(\hat{x})v\|$$

for every non-zero $v \in E^i_{\hat{x}}$ and $1 \le i \le k$. The dimension of $E^i_{\hat{x}}$ is called the *multiplicity* of $\lambda_i(\hat{x})$.

In this work we assume that the invariant measure $\hat{\mu}$ is ergodic. Then the Lyapunov exponents and the dimensions of the subspaces $E^i_{\hat{x}}$ are constant almost everywhere. The *Lyapunov spectrum* of the cocycle is the set of all Lyapunov exponents. The following notion is central to the whole paper: the Lyapunov spectrum is *simple* if it contains exactly d distinct Lyapunov exponents or, equivalently, if every Lyapunov exponent has multiplicity equal to 1.

2.2. **Partially hyperbolic skew-products.** Let $\hat{\sigma} : \hat{\Sigma} \to \hat{\Sigma}$ be any two-sided finite or countable shift. By this we mean that $\hat{\Sigma}$ is the set of two-sided sequences $(x_n)_{n \in \mathbb{Z}}$ in some set $X \subset \mathbb{N}$ with $\#X > 1$, and the map $\hat{\sigma}$ is given by

$$\hat{\sigma}\left((x_n)_{n \in \mathbb{Z}}\right) = (x_{n+1})_{n \in \mathbb{Z}}.$$

Let $\mathrm{dist}_{\hat{\Sigma}} : \hat{\Sigma} \times \hat{\Sigma} \to \mathbb{R}$ be the distance defined by

$$\mathrm{dist}_{\hat{\Sigma}}(\hat{x}, \hat{y}) = \sum_{k=-\infty}^{\infty} \frac{1}{2^{|k|}} \alpha_k \tag{1}$$

where $\hat{x} = (x_k)_{k \in \mathbb{Z}} \in \hat{\Sigma}$, $\hat{y} = (y_k)_{k \in \mathbb{Z}} \in \hat{\Sigma}$ and

$$\alpha_k = \begin{cases} 0 \text{ if } x_k = y_k \\ 1 \text{ otherwise.} \end{cases}$$

Then $\hat{\Sigma}$ is a metric space. Moreover, $\hat{\sigma}$ is a *hyperbolic homeomorphism* (in the sense of [19]), as we are going to explain.

Given any $\hat{x} \in \hat{\Sigma}$, we define the *local stable* and *unstable sets* of $\hat{x}$ with respect to $\hat{\sigma}$ by

$$W^s_{loc}(\hat{y}) = \{\hat{x} : x_k = y_k \text{ for every } k \geq 0\} \text{ and}$$
$$W^u_{loc}(\hat{y}) = \{\hat{x} : x_k = y_k \text{ for every } k \leq 0\}.$$

Observe that, taking $\lambda = 1/2$ and $\tau = 1/2$,

(i) $\operatorname{dist}(\hat{\sigma}^n(\hat{y}_1), \hat{\sigma}^n(\hat{y}_2)) \leq \lambda^n \operatorname{dist}(\hat{y}_1, \hat{y}_2)$ for any $\hat{y} \in \hat{\Sigma}$, $\hat{y}_1, \hat{y}_2 \in W^s_{loc}(\hat{y})$ and $n \geq 0$;

(ii) $\operatorname{dist}(\hat{\sigma}^{-n}(\hat{y}_1), \hat{\sigma}^{-n}(\hat{y}_2)) \leq \lambda^n \operatorname{dist}(\hat{y}_1, \hat{y}_2)$ for any $\hat{y} \in \hat{\Sigma}$, $\hat{y}_1, \hat{y}_2 \in W^u_{loc}(\hat{y})$ and $n \geq 0$;

(iii) if $\operatorname{dist}(\hat{x}, \hat{y}) \leq \tau$, then $W^s_{loc}(\hat{x})$ and $W^u_{loc}(\hat{y})$ intersect in a unique point, which is denoted by $[x, y]$ and depends continuously on $\hat{x}$ and $\hat{y}$.

By *partially hyperbolic skew-product* over the shift map $\hat{\sigma}$ we mean a homeomorphism $\hat{f} : \hat{\Sigma} \times K \to \hat{\Sigma} \times K$ of the form

$$\hat{f}(\hat{x}, t) = \big(\hat{\sigma}(\hat{x}), \hat{f}_{\hat{x}}(t)\big)$$

where K is a compact Riemann manifold and the maps $\hat{f}_{\hat{x}} : K \to K$ are diffeomorphisms satisfying

$$\lambda \|d\hat{f}_{\hat{x}}(t)\| < 1 \text{ and } \lambda \|d\hat{f}^{-1}_{\hat{x}}(t)\| < 1 \text{ for every } (\hat{x}, t) \in \hat{\Sigma} \times K, \tag{2}$$

where λ is a constant as in (i) - (ii). We also assume the following Hölder condition: there exist $C > 0$ and $\alpha > 0$ such that the C^1-distance between $\hat{f}_{\hat{x}}$ and $\hat{f}_{\hat{y}}$ is bounded by $C \operatorname{dist}(\hat{x}, \hat{y})^\alpha$ for every $\hat{x}, \hat{y} \in \hat{\Sigma}$.

We say that $\hat{f}$ has *mostly neutral center direction* if the family of maps $\hat{f}^n_{\hat{x}} : K \to K$ defined for $n \in \mathbb{Z}$ and $\hat{x} \in \hat{\Sigma}$ by

$$\hat{f}^n_{\hat{x}} = \begin{cases} \hat{f}_{\hat{\sigma}^{n-1}(\hat{x})} \circ \cdots \circ \hat{f}_{\hat{x}} & \text{if } n > 0 \\ \mathrm{id} & \text{if } n = 0 \\ \hat{f}^{-1}_{\hat{\sigma}^n(x)} \circ \cdots \circ \hat{f}^{-1}_{\hat{\sigma}^{-1}(\hat{x})} & \text{if } n < 0. \end{cases}$$

have bounded derivatives, that is, if there exists $C > 0$ such that

$$\|D\hat{f}^n_{\hat{x}}\| \leq C \text{ for every } \hat{x} \in \hat{\Sigma} \text{ and } n \in \mathbb{Z}.$$

Remark 2.1. Clearly, this implies that the $\{\hat{f}^n_{\hat{x}} : j \in \mathbb{Z} \text{ and } \hat{x} \in \hat{\Sigma}\}$ is equi-continuous. When the maps $\hat{f}_{\hat{y}}$ are $C^{1+\epsilon}$, equi-continuity alone suffices for all our purposes (see Remark 3.2).

A few comments are in order concerning the scope of the notion of partially hyperbolic skew-product. To begin with, the shift $\hat{\sigma} : \hat{\Sigma} \to \hat{\Sigma}$ may be replaced by a sub-shift $\hat{\sigma}_T : \hat{\Sigma}_T \to \hat{\Sigma}_T$ associated to a transition matrix $T = (T_{i,j})_{i,j \in X}$. By this we mean that $T_{i,j} \in \{0, 1\}$ for every $i, j \in X$ and $\hat{\sigma}_T$ is the restriction of the shift map $\hat{\sigma}$ to the subset $\hat{\Sigma}_T$ of sequences $(x_n)_{n \in \mathbb{Z}}$ such that $T_{x_n, x_{n+1}} = 1$ for every $n \in \mathbb{Z}$. One way to reduce the sub-shift case to the full shift case is through inducing. Namely, fix any 1-cylinder $[0; i] = \{(x_n)_{n \in \mathbb{Z}} \in \hat{\Sigma}_T : x_0 = i\}$ with positive measure and consider the first return map $g : [i] \to [i]$ of $\hat{\sigma}_T$ to $[i]$. This is conjugate to a full countable shift (with the return times as symbols) and it preserves the normalized restriction to the cylinder of the $\hat{\sigma}_T$-invariant measure. All the conditions that follow are not affected by this procedure. Moreover, every linear cocycle F over $\hat{\sigma}_T$ gives rise, also through inducing, to a linear cocycle over

g whose Lyapunov spectrum is just a rescaling of the Lyapunov spectrum of F. In particular, simplicity may also be read out from the induced cocycle.

Moreover, although we choose to formulate our approach in a symbolic set-up, for skew-products over shifts, it is clear that it extends to other situations that are more geometric in nature. For example, take $g : N \to N$ to be a partially hyperbolic diffeomorphism on a compact 3-dimensional manifold N and assume that there exists an embedded closed curve $\gamma \subset N$ such that $g(\gamma) = \gamma$ and some connected component of $W^s_{loc}(\gamma) \cap W^u_{loc}(\gamma) \setminus \gamma$ is a closed curve. By [8], g is conjugate up to finite covering to a skew-product over a linear Anosov diffeomorphism of the 2-torus. Thus, using a Markov partition for the Anosov map, one can semi-conjugate g to a partially hyperbolic skew-product over a sub-shift of finite type. In this way, the conclusions of this paper can be adapted to linear cocycles over such a diffeomorphism.

2.3. **Stable and unstable holonomies.** Property (2) is a condition of *domination* (or *normal hyperbolicity*, in the spirit of [16]): it means that any expansion and contraction of $\hat{f}_{\hat{x}}$ along the fibers $\{\hat{x}\} \times K$ are dominated by the hyperbolicity of the base map $\hat{\sigma}$. For our purposes, its main relevance is that it ensures the existence of strong-stable and strong-unstable "foliations" for $\hat{f}$, as we explain next.

Let the product $\hat{M} = \hat{\Sigma} \times K$ be endowed with the distance defined by

$$\mathrm{dist}_{\hat{M}}((\hat{x}_1, t_1), (\hat{x}_2, t_2)) = \mathrm{dist}_{\hat{\Sigma}}(\hat{x}_1, \hat{x}_2) + \mathrm{dist}_K(t_1, t_2),$$

where $\mathrm{dist}_{\hat{\Sigma}}$ denotes the distance (1) on $\hat{\Sigma}$ and dist_K is the distance induced by the Riemannian metric on K.

We consider the *stable holonomies*

$$h^s_{\hat{x},\hat{y}} : K \to K, \quad h^s_{\hat{x},\hat{y}} = \lim_{n\to\infty} \left(\hat{f}^n_{\hat{y}}\right)^{-1} \circ \hat{f}^n_{\hat{x}},$$

defined for every $\hat{x}$ and $\hat{y}$ with $\hat{x} \in W^s_{loc}(\hat{y})$, and *unstable holonomies*

$$h^u_{\hat{x},\hat{y}} : K \to K, \quad h^u_{\hat{x},\hat{y}} = \lim_{n\to-\infty} \left(\hat{f}^n_{\hat{y}}\right)^{-1} \circ \hat{f}^n_{\hat{x}}$$

defined for every $\hat{x}$ and $\hat{y}$ with $\hat{x} \in W^u_{loc}(\hat{y})$. That these families of maps exist follows from the assumption (2), using arguments from [6]; see for instance [5] which deals with a similar setting.

We define the *local strong-stable set* and the *local strong-unstable set* of each $(\hat{x}, t) \in \hat{M}$ to be

$$W^{ss}_{loc}(\hat{x}, t) = \{(\hat{y}, s) \in \hat{M} : \hat{y} \in W^s_{loc}(\hat{x}) \text{ and } s = h^s_{\hat{x},\hat{y}}(t)\} \text{ and}$$
$$W^{uu}_{loc}(\hat{x}, t) = \{(\hat{y}, s) \in \hat{M} : \hat{y} \in W^u_{loc}(\hat{x}) \text{ and } s = h^u_{\hat{x},\hat{y}}(t)\},$$

respectively. It is easy to check that

$$(\hat{y}, s) \in W^{ss}_{loc}(\hat{x}, t) \quad \Rightarrow \quad \lim_{n\to+\infty} \mathrm{dist}_{\hat{M}}(\hat{f}^n(\hat{y}, s), \hat{f}^n(\hat{x}, t)) = 0$$

and analogously on strong-unstable sets for time $n \to -\infty$.

2.4. **Measures with partial product structure.** Throughout, we take $\hat{\mu}$ to be an $\hat{f}$-invariant measure with *partial product structure*, that is, a probability measure of the form $\hat{\mu} = \hat{\rho}\, \mu^s \times \mu^u \times \mu^c$ where:

- $\hat{\rho} : \hat{M} \to (0, +\infty)$ is a continuous function bounded from zero and infinity;
- μ^s is a probability measure supported on $\Sigma^- = X^{\mathbb{Z}_{<0}}$;

- μ^u is a probability measure supported on $\Sigma^+ = X^{\mathbb{Z}_{\geq 0}}$;
- μ^c is a probability measure on the manifold K.

For notational convenience, we write the boundedness as follows: there exists $\kappa > 0$ such that

$$\frac{1}{\kappa} \leq \frac{\tilde\rho(x^s, x^u)}{\tilde\rho(x^s, z^u)} \leq \kappa \quad \text{and} \quad \frac{1}{\kappa} \leq \frac{\tilde\rho(x^s, x^u)}{\tilde\rho(z^s, x^u)} \leq \kappa \tag{3}$$

for every x^s, $z^s \in \Sigma^-$ and x^u, $z^u \in \Sigma^-$, where $\tilde\rho : \hat\Sigma \to \mathbb{R}$ is defined by

$$\tilde\rho(\hat x) = \int \hat\rho(\hat x, t) d\mu^c(t). \tag{4}$$

Observe that when $\hat\Sigma$ is a finite shift space this is an immediate consequence of compactness and the continuity of $\hat\rho$.

Now define, for $\hat x \in \hat M$,

$$\hat\varrho(\hat x, \cdot\,) = \frac{\hat\rho(\hat x, \cdot\,)}{\tilde\rho(\hat x)} \quad \text{and} \quad \hat\mu^c_{\hat x} = \hat\varrho(\hat x, \cdot\,)\, \mu^c. \tag{5}$$

In other words, $\hat\mu^c_{\hat x}$ is the normalization of $\hat\rho(\hat x, \cdot)\mu^c$. Note that the family $\{\hat\mu^c_{\hat x} : \hat x \in \hat\Sigma\}$ is a (continuous) disintegration of $\hat\mu$ along vertical fibers, that is, with respect to the partition $\hat{\mathcal{P}} = \{\{\hat x\} \times K : \hat x \in \hat\Sigma\}$.

The assumption that $\hat\mu$ is invariant under $\hat f$, together with the fact that $\hat\mu^c_{\hat x}$ depends continuously on $\hat x$, implies that

$$(\hat f_{\hat x})_* \hat\mu^c_{\hat x} = \hat\mu^c_{\hat\sigma(\hat x)} \quad \text{for every } \hat x \in \hat\Sigma. \tag{6}$$

We will also see in Section 5 that this disintegration is *holonomy invariant*:

$$\begin{aligned} &(h^s_{\hat x, \hat y})_* \hat\mu^c_{\hat x} = \hat\mu^c_{\hat y} \text{ whenever } \hat y \in W^s(\hat x) \quad \text{and} \\ &(h^u_{\hat x, \hat y})_* \hat\mu^c_{\hat x} = \hat\mu^c_{\hat y} \text{ whenever } \hat y \in W^u(\hat x). \end{aligned} \tag{7}$$

Remark 2.2. In particular, if $\hat p$ is a fixed point of the shift map then $\hat\mu^c_{\hat p}$ is invariant under $\hat f_{\hat p}$. Clearly, it is equivalent to μ^c. Moreover, if $\hat z$ is a homoclinic point of $\hat p$, that is, a point in $W^s(\hat p) \cap W^u(\hat p)$, then $(h^s_{\hat z, \hat p} \circ h^u_{\hat p, \hat z})_* \hat\mu^c_{\hat p} = (h^u_{\hat z, \hat p} \circ h^s_{\hat p, \hat z})_* \hat\mu^c_{\hat p} = \hat\mu^c_{\hat p}$.

2.5. **Linear cocycles with holonomies.** Let $\hat A : \hat M \to GL(d, \mathbb{C})$ be a α-Hölder continuous map for some $\alpha > 0$. By this we mean that there exists $C > 0$ such that

$$\|\hat A(\hat p) - \hat A(\hat q)\| \leq C \operatorname{dist}(\hat p, \hat q)^\alpha \quad \text{for any } \hat p, \hat q \in \hat M.$$

The linear cocycle defined by $\hat A$ over the transformation $\hat f : \hat M \to \hat M$ is the map $\hat F_{\hat A} : \hat M \times \mathbb{C}^d \to \hat M \times \mathbb{C}^d$ defined by

$$\hat F_{\hat A}(\hat p, v) = \big(\hat f(\hat p), \hat A(\hat p) v\big).$$

In what follows we take the cocycle to admit *strong-stable holonomies* and *strong-unstable holonomies*, in the sense of [6, 2]. Let us explain this.

By *strong-stable holonomies* we mean a family of linear transformations $H^s_{\hat p, \hat q} : \mathbb{C}^d \to \mathbb{C}^d$, defined for each $\hat p, \hat q \in \hat M$ with $\hat q \in W^{ss}_{loc}(\hat p)$ and such that, for some constant $L > 0$,

(a) $H^s_{\hat f^j(\hat p), \hat f^j(\hat q)} = \hat A^j(\hat q) \circ H^s_{\hat p, \hat q} \circ \hat A^j(\hat p)^{-1}$ for every $j \geq 1$;
(b) $H^s_{\hat p, \hat p} = \mathrm{id}$ and $H^s_{\hat p, \hat q} = H^s_{\hat z, \hat q} \circ H^s_{\hat p, \hat z}$ for any $\hat z \in W^{ss}_{loc}(\hat p)$;
(c) $\|H^s_{\hat p, \hat q} - \mathrm{id}\,\| \leq L \operatorname{dist}(\hat p, \hat q)^\alpha$.

Strong unstable holonomies $H^u_{\hat{p},\hat{q}} : \mathbb{C}^d \to \mathbb{C}^d$ are defined analogously, for the pairs $(\hat{p}, \hat{q})$ with $\hat{q} \in W^{uu}_{loc}(\hat{p})$.

It was shown in [2] that strong-stable holonomies and strong-unstable holonomies do exist, in particular, when the cocycle is *fiber bunched*. By the latter we mean that there exist $C > 0$ and $\theta < 1$ such that

$$\|\hat{A}^n(\hat{p})\|\|\hat{A}^n(\hat{p})^{-1}\|\lambda^{n\alpha} \le C\theta^n \quad \text{for every } \hat{p} \in \hat{M} \text{ and } n \ge 0,$$

where λ is a hyperbolicity constant for $\hat{f}$ as in conditions (i)-(ii) above.

For $1 \le l \le d-1$, let $\mathrm{Sec}(K, \mathrm{Grass}(l,d))$ denote the space of measurable maps V from (some full μ^c-measure subset of) K to the Grassmannian manifold of all l-dimensional subspaces of $\mathbb{R}^d$. For each $\hat{x} \in \hat{\Sigma}$, consider the following push-forward maps

$$\mathrm{Sec}(K, \mathrm{Grass}(l,d)) \to \mathrm{Sec}(K, \mathrm{Grass}(l,d)) :$$

(a) $V \mapsto \mathcal{F}_{\hat{x}} V$ given by

$$\mathcal{F}_{\hat{x}} V(t) = \hat{A}(\hat{x}, s)V(s) \text{ with } s = (\hat{f}_{\hat{x}})^{-1}(t);$$

(b) $V \mapsto \mathcal{H}^s_{\hat{x},\hat{y}} V$ given, for $\hat{y} \in W^s_{loc}(\hat{x})$, by

$$\mathcal{H}^s_{\hat{x},\hat{y}} V(t) = H^s_{(\hat{x},s),(\hat{y},t)} V(s) \text{ with } s = h^s_{\hat{y},\hat{x}}(t);$$

(c) $V \mapsto \mathcal{H}^u_{\hat{x},\hat{y}} V$ given, for $\hat{y} \in W^u_{loc}(\hat{x})$, by

$$\mathcal{H}^u_{\hat{x},\hat{y}} V(t) = H^u_{(\hat{x},s),(\hat{y},t)} V(s) \text{ with } s = h^u_{\hat{y},\hat{x}}(t).$$

2.6. **Pinching and twisting.** Now we state our refined criterion for simplicity of the Lyapunov spectrum.

We call the cocycle $\hat{F}$ *pinching* if there exists some fixed (or periodic) vertical leaf $\ell = \{\hat{p}\} \times K$ such that the restriction to ℓ of every exterior power $\Lambda^k \hat{F}$ has simple Lyapunov spectrum, relative to the $\hat{f}_{\hat{p}}$-invariant measure $\hat{\mu}^c_{\hat{p}}$ (Remark 2.2). In other words, the Lyapunov exponents $\lambda_1, \cdots, \lambda_d$ are such that, for each $1 \le k \le d-1$ and $\hat{\mu}^c_{\hat{p}}$-almost every $t \in K$, the sums

$$\lambda_{i_1}(\hat{p}, t) + \cdots + \lambda_{i_k}(\hat{p}, t), \qquad 1 \le i_1 < \cdots < i_k \le d$$

are all distinct. It is clear that $\hat{F}$ is pinching if it is uniformly pinching.

Take $\hat{F}$ to be pinching and let $E^1(t) \oplus \cdots \oplus E^d(t)$ denote the Oseledets decomposition of $\hat{F}$ restricted to ℓ. The maps $t \mapsto E^i(t)$ are defined on a full $\hat{\mu}^c_{\hat{p}}$-measure set. Now let $\hat{z}$ be some homoclinic point of $\hat{p}$ and $\imath \ge 1$, such that

$$\hat{z} \in W^u_{loc}(\hat{p}) \quad \text{and} \quad \hat{\sigma}^\imath(\hat{z}) \in W^s_{loc}(\hat{p}).$$

Define

$$V^i = (\mathcal{H}^u_{\hat{z},\hat{p}} \circ \mathcal{F}^{-\imath}_{\hat{z}} \circ \mathcal{H}^s_{\hat{p},\hat{\sigma}^\imath(\hat{z})}) E^i \quad \text{for } i = 1, \ldots, d. \tag{8}$$

Note that V^i is also defined on a full $\hat{\mu}^c_{\hat{p}}$-measure set since the maps $\hat{f}_{\hat{x}}$ and the holonomies $h^s_{\hat{x},\hat{y}}$ and $h^u_{\hat{x},\hat{y}}$ preserve $\hat{\mu}^c_{\hat{p}}$ (Remark 2.2).

Now let $B(t) = (\beta_{i,j}(t))_{i,j}$ be the $d \times d$-matrix defined by

$$V^i(t) = \sum_{j=1}^{d} \beta_{i,j}(t) E^j(t), \quad \text{for } i = 1, \ldots, d.$$

We call the cocycle $\hat{F}$ *twisting* if, some choice of the homoclinic point $\hat{z}$, all the algebraic minors $m_{I,J}(t)$ of $B(t)$ decay sub-exponentially along the orbits of $\hat{f}_{\hat{p}}$, meaning that

$$\lim_{n\to\infty} \frac{1}{n} \log |m_{I,J}(\hat{f}^n_{\hat{p}}(t))| = 0 \quad \text{for } \hat{\mu}^c_{\hat{p}}\text{-almost every } t \in K \tag{9}$$

and every $I, J \subset \{1, \dots, d\}$ with $\#I = \#J$.

It is clear that this holds if $\hat{F}$ is uniformly twisting, because in this case the algebraic minors are uniformly bounded. More generally (see for instance [20, Corollary 3.11]), the property (9) holds whenever the function $\log |m_{I,J}| \circ \hat{f}_{\hat{p}} - \log |m_{I,J}|$ is $\hat{\mu}^c_{\hat{p}}$-integrable.

2.7. **Main statements and outline of the proof.** We say that the cocycle $\hat{F}$ is *simple* if it is both pinching and twisting, in the sense of the previous section. That is the case, in particular, if $\hat{F}$ is uniformly pinching and uniformly twisting. Thus Theorem A is contained in the following result:

Theorem B. If $\hat{F} : \hat{M} \times \mathbb{C}^d \to \hat{M} \times \mathbb{C}^d$ is a simple cocycle then its Lyapunov spectrum is simple.

Let $H^\alpha(\hat{M})$ denote the space of all α-Hölder continuous maps $\hat{A} : \hat{M} \to GL(d, \mathbb{C})$. The norm

$$\|\hat{A}\|_\alpha = \sup_{\hat{x}\in\hat{M}} \|\hat{A}(\hat{x})\| + \sup_{\hat{x}\neq\hat{y}} \frac{\|\hat{A}(\hat{x}) - \hat{A}(\hat{y})\|}{\operatorname{dist}(\hat{x}, \hat{y})^\alpha}$$

defines a topology in $H^\alpha(\hat{M})$ that we call *α-Hölder topology*. We are going to prove in Section 10 that the uniform pinching and uniform twisting conditions are open with respect to this topology. Thus Theorem A implies

Theorem C. There is a non-empty open subset of $\hat{A} \in H^\alpha(\hat{M})$ such that the associated linear cocycle $\hat{F}$ over $\hat{f}$ has simple Lyapunov spectrum.

In many contexts of linear cocycles over *hyperbolic* systems, simplicity turns out to be a generic condition: it contains an open and dense subset of cocycles (precise statements can be found in Viana [20]). This is related to the fact that, in the hyperbolic setting, the pinching and twisting conditions are just transversality conditions, they clearly hold on the complement of suitable submanifolds with positive codimension. At present, it is unclear how this can be extended to the partially hyperbolic setting. Uniform pinching is surely not a generic condition, in general, but it might be locally generic in some special situations, for instance when $\hat{f}_{\hat{p}}$ is quasi-periodic.

We close this section by outlining the overall strategy of the proof of Theorem B. For every $1 \le \ell < d$, we want to find complementary $\hat{F}-$invariant measurable sections

$$\xi : \hat{M} \to \operatorname{Grass}(l, d) \quad \text{and} \quad \eta : \hat{M} \to \operatorname{Grass}(d - l, d) \tag{10}$$

such that the Lyapunov exponents of $\hat{F}$ along ξ are strictly larger than those along η.

The starting point is to reduce the problem to the case when the maps $\hat{f}_{\hat{x}}$ and the matrices $\hat{A}(\hat{x}, t)$ depend on $\hat{x}$ only through its positive part x^u. This we do in Section 4, using the stable holonomies to conjugate the original dynamics to others with these properties. Then $\hat{f} : \hat{M} \to \hat{M}$ projects to a transformation $f : M \to M$

on $M = \Sigma^+ \times K$ which is a skew-product over the one-sided shift $\sigma : \Sigma^+ \to \Sigma^+$ and, similarly, the linear cocycle $\hat{F} : \hat{M} \times \mathbb{C}^d \to \hat{M} \times \mathbb{C}^d$ projects to a linear cocycle $F : M \times \mathbb{C}^d \to M \times \mathbb{C}^d$ over the transformation $\hat{f}$.

We also denote by $\hat{F}$ and F the actions

$$\hat{F} : \hat{M} \times \mathrm{Grass}(l,d) \to \hat{M} \times \mathrm{Grass}(l,d) \text{ and}$$
$$F : M \times \mathrm{Grass}(l,d) \to M \times \mathrm{Grass}(l,d)$$

induced by the two linear cocycles on the Grassmannian bundles. Still in Section 4, using very classical arguments, we relate the invariant measures of $\hat{f}$ and $\hat{F}$ with those of f and F, respectively.

In Section 5 we study *u-states*, that is $\hat{F}$-invariant probability measures $\hat{m}$ whose Rokhling disintegrations $\{\hat{m}_{\hat{x}} : \hat{x} \in \hat{\Sigma}\}$ are invariant under unstable holonomies, as well as the corresponding F-invariant probability measures m. Here we meet the first important new difficulty arising from the fact that $\hat{f}$ is only partially hyperbolic.

Indeed, in the hyperbolic setting such measures m are known to admit *continuous* disintegrations $\{m_x : x \in M\}$ along the fibers $\{x\} \times \mathrm{Grass}(l,d)$ and this fact plays a key part in the arguments of Bonatti-Viana [7] and Avila-Viana [3].

In the partially hyperbolic setting, the situation is far more subtle: the disintegration $\{m_x : x \in \Sigma\}$ along the sets $\{x\} \times K \times \mathrm{Grass}(l,d)$ is still continuous, but there is no reason why this should extend to the disintegration

$$\{m_{x,t} : (x,t) \in M\}$$

along fibers $K \times \mathrm{Grass}(l,d)$, which is what one really needs. The way we make up for this is by proving a kind of L^1-continuity: if $(x_i)_i \to x$ in Σ then $(m_{x_i,t})_i \to m_{x,t}$ in $L^1(\mu^c)$. See Proposition 5.9 for the precise statement.

This also leads to our formulating the arguments in terms of measurable sections $K \to \mathrm{Grass}(l,d)$ of the Grassmannian bundle, which is perhaps another significant novelty in this paper. The properties of such sections are studied in Section 6. The key result (Proposition 6.1) is that, under pinching and twisting, the graph of every invariant Grassmannian section has zero m_x-measure, for every $x \in M$ and any u-state $\hat{m}$.

These results build up to Section 7, where we prove that every u-state $\hat{m}$ has an atomic disintegration. More precisely (Theorem 7.1), there exists a measurable section $\xi : \hat{M} \to \mathrm{Grass}(l,d)$ such that, given any u-state $\hat{m}$ on $\hat{M} \times \mathrm{Grass}(l,d)$, we have

$$\hat{m}_{\hat{x},t} = \delta_{\xi(\hat{x},t)} \quad \text{for } \hat{\mu}\text{-almost every } (\hat{x},t) \in \hat{M}. \tag{11}$$

Thus we construct the invariant section $\xi : \hat{M} \to \mathrm{Grass}(l,d)$ in (10).

To find the complementary invariant section $\eta : \hat{M} \to \mathrm{Grass}(d-l,d)$, in Section 8 we apply the same procedure to the adjoint cocycle $\hat{F}^*$, that is, the linear cocycle defined over $\hat{f}^{-1} : \hat{M} \to \hat{M}$ by the function

$$\hat{x} \mapsto \hat{A}^*(x) = \text{ adjoint of } \hat{A}(\hat{f}^{-1}(\hat{x})).$$

We check (Proposition 8.4) that this cocycle $\hat{F}^*$ is pinching and twisting if and only if $\hat{F}$ is. So, the previous arguments yield a $\hat{F}^*$-invariant section $\xi^* : \hat{M} \to \mathrm{Grass}(l,d)$ related to the u-states of $\hat{F}^*$. Then we just take $\eta = \left(\xi^*\right)^\perp$.

Finally, in Section 9 we check that the eccentricity, or lack of conformality, of the iterates $\hat{A}^n$ goes to infinity $\hat{\mu}$-almost everywhere (see Proposition 9.1) and we use this fact to deduce that every Lyapunov exponent of $\hat{F}$ along ξ is strictly larger than any of the Lyapunov exponents of $\hat{F}$ along η. At this stage the arguments are again very classical. This concludes the proof of Theorem B.

Theorem C is proven in Section 10. The two appendices contain material that seems to be folklore, but for which we could not find explicit references. In Appendix A we show that continuous maps are dense in the corresponding L^1 space, whenever the target space is geodesically convex.

3. Disintegration along center leaves

Let us start by fixing some terminology, once and for all. We use id_Y to denote the identity transformation in a set Y. Similarly, dist_Y will always denote the distance in a metric space Y.

Recall that $\hat{\Sigma} = \Sigma^- \times \Sigma^+$. Thus we write every $\hat{x} \in \hat{\Sigma}$ as (x^s, x^u) with $x^s \in \Sigma^-$ and $x^u \in \Sigma^+$. For simplicity, we also write $\Sigma = \Sigma^+$ and $x = x^u$. Let $P : \hat{\Sigma} \to \Sigma$ be the canonical projection given by $P(\hat{x}) = x$ and let $\sigma : \Sigma \to \Sigma$ be the one-sided shift.

Given points $\hat{x} \in \hat{\Sigma}$ and $\hat{q} \in \hat{M}$, denote

$$x_n = P(\hat{\sigma}^{-n}(\hat{x})) \quad \text{and} \quad q_n = (P \times \mathrm{id}_K)(\hat{f}^{-n}(\hat{q}))$$

for each $n \in \mathbb{N}$. In what follows, $\hat{p} \in \hat{M}$ is a fixed point of $\hat{\sigma}$ and $\hat{z} \in \hat{M}$ is a homoclinic point of $\hat{p}$. More precisely, we fix $\hat{z}$ and some $\imath \in \mathbb{N}$ so that

$$\hat{z} \in W^u_{loc}(\hat{p}) \quad \text{and} \quad \hat{\sigma}^\imath(\hat{z}) \in W^s_{loc}(\hat{p}).$$

We also consider $M = \Sigma \times K$ and the projection $\mu = (P \times \mathrm{id}_K)_* \hat{\mu}$. In other words,

$$\mu = \rho(x,t)\, \mu^u \times \mu^c \quad \text{where} \quad \rho(x,t) = \int \hat{\rho}(x^s, x, t)\, d\mu^s(x^s). \tag{12}$$

Similarly to (5), for each $x \in M$ let

$$\varrho(x,\cdot) = \frac{\rho(x,\cdot)}{\int \rho(x,t)\, d\mu^c(t)} \quad \text{and} \quad \mu^c_x = \varrho(x,\cdot)\mu^c. \tag{13}$$

Note that $\{\mu^c_x : x \in \Sigma\}$ is a continuous disintegration of μ with respect to the partition $\mathcal{P} = \{\{x\} \times K : x \in \Sigma\}$.

In this section we derive some useful properties of these disintegrations (5) and (13). For this, we assume that the base dynamics is such that each $\hat{f}_{\hat{x}} : K \to K$ along the center direction depends only on $x = P(\hat{x})$. This is no restriction in our setting, as we will see in Section 4. Then there exists $f : M \to M$ of the form $f(x,t) = (\sigma(x), f_x(t))$ such that

$$(P \times \mathrm{id}_K) \circ \hat{f} = f \circ (P \times \mathrm{id}_K)$$

3.1. **Holonomy invariance.** We call the *extremal center Lyapunov exponents* of $\hat{f}$ the limits

$$\lambda^{c+} = \lim_n \frac{1}{n} \log \left\| D\hat{f}^n_{\hat{x}}(t) \right\| \quad \text{and} \quad \lambda^{c-} = \lim_n -\frac{1}{n} \log \left\| D\hat{f}^n_{\hat{x}}(t)^{-1} \right\|.$$

The Oseledets theorem [17] ensures that these numbers are well defined at $\hat{\mu}$-almost every point. In our situation:

Lemma 3.1. $\lambda^{c+} = \lambda^{c-} = 0$.

Proof. A trivial consequence of the assumption that the maps $\hat{f}^n_{\hat{x}}$ have uniformly bounded derivatives. □

Remark 3.2. When the maps $\hat{f}^n_{\hat{x}}$ are $C^{1+\epsilon}$, equi-continuity alone suffices to get the conclusion of Lemma 3.1. This can be shown using Pesin theory, as follows.

Suppose that $\lambda^{c+} > 0$. Then we have a Pesin unstable manifold defined $\hat{\mu}$-almost everywhere. This implies that there exist $\hat{x} \in \hat{\Sigma}$ and $t \neq s \in K$ such that

$$\mathrm{dist}_K\left(\hat{f}^{-n}_{\hat{x}}(t), \hat{f}^{-n}_{\hat{x}}(s)\right) \to 0.$$

Then, given points t and s in the unstable manifold and given any $\delta > 0$, there exists n such that $\mathrm{dist}_K(\hat{f}^{-n}_{\hat{x}}(t), \hat{f}^{-n}_{\hat{x}}(s)) < \delta$. This implies that the family is not equi-continuous. The proof for λ^{c-} is analogous.

Let $\pi_1 : \hat{M} \to \hat{\Sigma}$ be the projection $\pi_1(\hat{x}, t) = \hat{x}$, recall that we also assume that $\hat{f} : \hat{M} \to \hat{M}$ admits s-holonomies and u-holonomies and $\hat{\mu}$ has partial product structure. That implies that $(\pi_1)_*\hat{\mu}$ has local product structure in the sense of [4].

Lemma 3.3. *The map $\hat{x} \mapsto \hat{\mu}^c_{\hat{x}}$ is continuous. Moreover, the disintegration $\{\hat{\mu}^c_{\hat{x}} : \hat{x} \in \hat{M}\}$ is both u-invariant and s-invariant:*

(a) $\left(h^u_{\hat{x},\hat{y}}\right)_* \hat{\mu}^c_{\hat{x}} = \hat{\mu}^c_{\hat{y}}$ *for every* $\hat{x} \in W^u(\hat{y})$ *and*

(b) $\left(h^s_{\hat{x},\hat{z}}\right)_* \hat{\mu}^c_{\hat{x}} = \hat{\mu}^c_{\hat{z}}$ *for every* $\hat{x} \in W^s(\hat{z})$.

Proof. By Theorem D in [4], there exists a disintegration $\{\tilde{\mu}^c_{\hat{x}} : \hat{x} \in \hat{\Sigma}\}$ which is continuous, u-invariant and s-invariant. By essential uniqueness, $\tilde{\mu}^c_{\hat{x}} = \hat{\mu}^c_{\hat{x}}$ for $\hat{\mu}$-almost every x. Since both disintegrations are continuous, it follows that they coincide, and so $\{\hat{\mu}^c_{\hat{x}} : \hat{x} \in \hat{\Sigma}\}$ is continuous, u-invariant and s-invariant, as claimed. □

Corollary 3.4. $\mu^c_x = \hat{\mu}^c_{\hat{x}}$ *for every* $\hat{x} \in \hat{\Sigma}$, *where* $x = P(\hat{x})$.

Proof. The assumption that $\hat{f}_{\hat{x}}$ only depends on $x = P(\hat{x})$ implies that $h^s_{\hat{x},\hat{y}} = \mathrm{id}_K$ for every $\hat{x}$ and $\hat{y}$ in the same stable set. By the previous lemma, this implies that $\hat{\mu}^c_{\hat{x}} = \hat{\mu}^c_{\hat{y}}$ whenever $\hat{x}$ and $\hat{y}$ are in the same stable set. Then,

$$\mu^c_x = \int \hat{\mu}^c_{\hat{y}}\, d\hat{\mu}^s_x(\hat{y}) = \hat{\mu}^c_{\hat{x}}$$

for any $\hat{x}$ with $P(\hat{x}) = x$. □

We also have

Corollary 3.5. *The disintegration $\{\mu^c_x : x \in \Sigma\}$ is f-invariant, in the sense that $\left(f_x\right)_* \mu^c_x = \mu_{\sigma(x)}$ for every $x \in \Sigma$.*

Proof. We have that $(\hat{f}_{\hat{x}})_*\hat{\mu}^c_{\hat{x}} = \hat{\mu}^c_{\hat{\sigma}(\hat{x})}$ for $\hat{\mu}$-almost every $\hat{x}$, because $\hat{\mu}$ is $\hat{f}$-invariant. Since $\hat{x} \mapsto \hat{\mu}^c_{\hat{x}}$ is continuous, the identity extends to *every* $\hat{x} \in \hat{\Sigma}$. By Corollary 3.4 this implies that $(f_x)_*\mu^c_x = \mu^c_{\sigma(x)}$ for every $x \in \Sigma$. □

3.2. **Jacobians.** Denote $\hat\nu = (\pi_1)_*\hat\mu$ and $\nu = (\pi_1)_*\mu$ where π_1 denotes both canonical projections $\hat M \to \hat\Sigma$ and $M \to \Sigma$. Recall the functions $\hat\varrho$ and ϱ defined in (5) and (13). Note that $\{\hat\varrho(x)\hat\mu^c : \hat x \in \hat\Sigma\}$ is a disintegration of $\hat\mu$ with respect to the partition $\{\pi_1^{-1}(\hat x) : \hat x \in \hat\Sigma\}$ of $\hat M$ and $\{\varrho(x)\mu^c : x \in \Sigma\}$ is a disintegration of μ with respect to the partition $\{\pi_1^{-1}(x) : x \in \Sigma\}$ of $\hat M$.

Remark 3.6. Let $Jf_x^j : K \to \mathbb{R}$ be the Jacobian of f_x^j with respect to μ^c. Using the observation that $\{\varrho(x)\mu^c : x \in \Sigma\}$ is a disintegration of the f-invariant measure μ, one easily gets that

$$Jf_x^j(t) = \frac{\varrho(f^j(x,t))}{\varrho(x,t)}.$$

In particular, these Jacobians are uniformly bounded from above and below. Analogously, the fact that $\{\hat\mu_{\hat x}^c : \hat x \in \hat\Sigma\}$ is invariant under s-holonomies and u-holonomies ensures that the Jacobians $Jh^*_{\hat x,\hat y}$ of those holonomies with respect to μ^c are uniformly bounded from above and below.

Lemma 3.7. *$J_\mu f^k(x,t) = J_\nu\sigma^k(x)$ for every $(x,t) \in M$ and $k \ge 1$.*

Proof. Fix some k-cylinder $I = [0; x_0, \dots, x_{k-1}]$ and let $J \subset I$ and $C \subset K$. Noting that $\sigma^k \mid I$ is injective,

$$\begin{aligned}\mu\big(f^k(J\times C)\big) &= \int_{y\in\sigma^k(J)} \mu_y^c(f^k_{z(y)}(C))\, d\nu(y)\\ &= \int_{z\in J} J_\nu\sigma^k(z)\mu^c_{\sigma^k(z)}(f^k_z(C))\, d\nu(z),\end{aligned}$$

where $z(y)$ is the unique point in $\sigma^{-k}(y)\cap I$ and we use the change of variables $z = z(y)$. Using Lemma 3.5, it follows that

$$\mu\big(f^k(J\times C)\big) = \int_J J_\nu\sigma^k(z)\mu_z^c(C)\, d\nu(z) = \int_{J\times C} (J_\nu\sigma^k\circ\pi_1)\, d\mu,$$

which concludes the proof. □

Now we find the Jacobian of σ^k:

Lemma 3.8. *$J_\nu\sigma^k(x) = 1/\nu_x^s(I)$, where $I = [-k; x_0, \dots, x_{k-1}]$. Consequently, the Jacobians $J_\mu f^k = J_\nu\sigma^k \circ \pi_1$ are continuous and bounded from zero and infinity on every k-cylinder.*

Proof. Given $x \in \Sigma$ and $n \ge 1$, let $J_n = [x_0, \dots, x_n]$ be the n-cylinder that contains x. Then,

$$J_\nu\sigma^k(x) = \lim_{n\to\infty} \frac{\nu(\sigma^k(J_n))}{\nu(J_n)}$$

Since $\hat\nu$ is invariant under $\hat\sigma$,

$$\nu(J_n) = \hat\nu(\Sigma^-\times J_n) = \hat\nu(\hat\sigma^k(\Sigma^-\times J_n)) = \int_{y\in\sigma^k(J_n)} \nu_y^s(I) d\nu(y).$$

It follows that

$$\frac{1}{J_\nu\sigma^k(x)} = \lim_{n\to\infty} \frac{\int_{y\in\sigma^k(J_n)} \nu_y^s(I) d\nu(y)}{\nu(\sigma^k(J_n))} = \nu_x^s(I).$$

This proves the first part of the conclusion. The second part is a consequence, since the local product structure implies that $x \mapsto \nu_x^s(I)$ is continuous for every cylinder I. □

4. Convergence of conditional measures

For each $1 \le l < d$, the linear cocycle $\hat{F} : \hat{M} \times \mathbb{C}^d \to \hat{M} \times \mathbb{C}^d$ induces a projective cocycle $\hat{F} : \hat{M} \times \mathrm{Grass}(l,d) \to \hat{M} \times \mathrm{Grass}(l,d)$ through

$$\hat{F}(\hat{q}, v) = (\hat{f}(\hat{q}), \hat{A}(\hat{q})v). \tag{14}$$

Denote $\mu = (P \times \mathrm{id}_K)_* \, \hat{\mu}$ and, given any Borel probability measure $\hat{m}$ on $\hat{M} \times \mathrm{Grass}(l,d)$,

$$m = \left(P \times \mathrm{id}_K \times \mathrm{id}_{\mathrm{Grass}(l,d)}\right)_* \hat{m}. \tag{15}$$

We will be especially interested in the case when $\hat{m}$ is a $\hat{f}$-invariant probability measure that projects down to $\hat{\mu}$ under the canonical projection $\pi : \hat{M} \times \mathrm{Grass}(l,d) \to \hat{M}$ on the first coordinate.

4.1. **Reduction to the one-sided case.** Our first step is to show that, up to conjugating the cocycle in a suitable way, we may suppose that:

(A) the base dynamics $\hat{f}_{\hat{x}}$ along the center direction depends only on x;
(B) the matrix $\hat{A}(\hat{x}, t)$ depend only on (x, t).

Let us explain how such a conjugacy may be defined using the stable holonomies.

Let $x^s \in \Sigma^-$ be fixed. For any $\hat{y} \in \hat{\Sigma}$, let $\phi(\hat{y}) = (x^s, y)$ and then define

$$h(\hat{y}, t) = \left(\hat{y}, h^s_{\varphi(\hat{y}), \hat{y}}(t)\right).$$

Then $\tilde{f} = h^{-1} \circ f \circ h$ is given by

$$\tilde{f}(\hat{y}, t) = \left(\hat{\sigma}(\hat{y}), \tilde{f}_{\hat{y}}(t)\right), \quad \text{with } \tilde{f}_{\hat{y}}(t) = h^s_{\hat{\sigma}(\phi(\hat{y})), \phi(\hat{\sigma}(\hat{y}))} f_{\phi(\hat{y})}(t).$$

Notice that $\tilde{f}_{\hat{y}}$ does depend only on y (because ϕ does).

Assume that (A) is satisfied. Define $\hat{\phi}(\hat{y}, t) = (\phi(\hat{y}), t)$ and then let

$$H(\hat{y}, t) = H^s_{\hat{\phi}(\hat{y}, t), (\hat{y}, t)}$$

Define $\tilde{A}(\hat{y}, t) = H(\hat{f}(\hat{y}, t))^{-1} \circ A(\hat{y}, t) \circ H(\hat{y}, t)$. Then

$$\tilde{A}(\hat{y}, t) = H^s_{\hat{f}(\hat{\phi}(\hat{y}, t)), \hat{\phi}(\hat{f}(\hat{y}, t))} \circ \hat{A}(\hat{\phi}(\hat{y}, t)),$$

which only depends on (y, t). Clearly, this procedure does not affect the Lyapunov exponents.

From now on, we assume that both (A) and (B) are satisfied. Then, there exist

$$f : M \to M, \ f(x, t) = (\sigma(x), f_x(t)) \quad \text{and} \quad A : M \to GL(d, \mathbb{C})$$

such that

$$(P \times \mathrm{id}_K) \circ \hat{f} = f \circ (P \times \mathrm{id}_K) \quad \text{and} \quad \hat{A} = A \circ (P \times \mathrm{id}_K).$$

Consequently, the map

$$F : M \times \mathrm{Grass}(l,d) \to M \times \mathrm{Grass}(l,d), \quad F(p, V) = (f(p), A(p)V)$$

satisfies

$$\left(P \times \mathrm{id}_K \times \mathrm{id}_{\mathrm{Grass}(l,d)}\right) \circ \hat{F} = F \circ \left(P \times \mathrm{id}_K \times \mathrm{id}_{\mathrm{Grass}(l,d)}\right).$$

The following general fact will be used to characterize the F-invariant probability measures:

Proposition 4.1. *Let $(N, \mathfrak{B}, \eta)$ be a Lebesgue probability space and $g : N \to N$ be a measurable map that preserves η. Let $\{\eta_y : y \in N\}$ be the disintegration of η with respect to the partition into pre-images $\mathcal{P} = \{g^{-1}(y) : y \in N\}$. Let*

$$G : N \times L \to N \times L, \quad G(x, v) = (g(x), G_x(v))$$

be a measurable skew-product over g and, given any probability measure m on $N \times L$ that projects down to η, let $\{m_x : x \in N\}$ be its disintegration with respect to the partition into vertical fibers $\{x\} \times L$, $x \in N$. Then m is invariant under G if and only if

$$m_x = \int (G_z)_* m_z \, d\eta_x(z) \quad \text{for } \eta\text{-almost every } x \in N.$$

Proof. Let us begin with some explanations concerning our notations. Let $\pi : N \to \mathcal{P}$ be the canonical quotient map, assigning to each $x \in N$ the element $\pi(x)$ of $\mathcal{P}$ that contains x. By definition, the Rokhlin disintegration of η with respect to $\mathcal{P}$ is the (essentially unique) measurable family $\{\eta_P : P \in \mathcal{P}\}$ of probability measures such that $\eta_P(P) = 1$ for each $P \in \mathcal{P}$ and

$$\begin{aligned} \int \phi(z) \, d\eta(z) &= \int \int \phi(z) \, d\eta_P(z) \, d(\pi_* \eta)(P) \\ &= \int \int \phi(z) \, d\eta_{\pi(x)}(z) \, d\eta(x). \end{aligned} \tag{16}$$

for any bounded measurable function $\phi : N \to \mathbb{R}$. Let $p : N \to \mathcal{P}$ be defined by $p(y) = g^{-1}(y)$. Observe that p is a bijection; thus we simply write $\eta_{p(y)} = \eta_y$ for each $y \in N$. Observe also that

$$\pi(x) = g^{-1}(g(x)) = p(g(x)) \text{ for every } x \in N.$$

Thus, (16) may be rewritten as

$$\begin{aligned} \int \phi(z) \, d\eta(z) &= \int \int \phi(z) \, d\eta_{\pi(g(x))} \, d\eta(x) \\ &= \int \int \phi(z) \, d\eta_{g(x)} \, d\eta(x) \end{aligned} \tag{17}$$

for any bounded measurable function $\phi : N \to \mathbb{R}$.

Define $\tilde{m}_x = \int (G_z)_* m_z \, d\eta_x(z)$ for each $x \in N$. Start by observing that

$$\begin{aligned} \tilde{m}_x\big(\{x\} \times L\big) &= \int m_z\big(\{z\} \times G_z^{-1}(L)\big) \, d\eta_x(z) \\ &= \int m_z\big(\{z\} \times L\big) \, d\eta_x(z) = 1 \end{aligned}$$

for each $x \in N$. Next, consider any measurable set $A \subset N \times L$. By the definition and the assumption that η is invariant under g,

$$\begin{aligned} \int \tilde{m}_x(A) \, d\eta(x) &= \int \int m_z\big(G_z^{-1}(A)\big) \, d\eta_x(z) \, d\eta(x) \\ &= \int \int m_z\big(G_z^{-1}(A)\big) \, d\eta_{g(x)}(z) \, d\eta(x). \end{aligned}$$

By (17), this means that

$$\int \tilde{m}_x(A) \, d\eta(x) = \int m_z(G^{-1}(A)) \, d\eta(z) = m(G^{-1}(A)) = G_* m(A)$$

for any measurable set $A \subset N \times L$.

This proves that $\{\tilde{m}_x : x \in N\}$ is a disintegration of G_*m with respect to the partition into vertical fibers. By the essential uniqueness of the disintegration, it follows that $m = G_*m$ if and only if $m_x = \tilde{m}_x$ for η-almost every x. □

As an immediate consequence, we get:

Corollary 4.2. *In the conditions of Proposition 4.1, if g is invertible then m is invariant under G if and only if $m_x = (G_{g^{-1}(x)})_* m_{g^{-1}(x)}$ for η-almost every $x \in N$.*

Proof. Each η_y must coincide with the Dirac mass at $g^{-1}(y)$. □

4.2. **Lifting of measures.** The next proposition shows that every $\hat{F}$-invariant measure $\hat{m}$ that projects down to $\hat{m}$ may be recovered from the corresponding F-invariant measure m, defined by (15). Recall that we write $q_n = (P \times \mathrm{id}_K)(\hat{f}^{-n}(\hat{q}))$ for each $\hat{q} \in \hat{M}$ and $n \geq 0$.

Proposition 4.3. *Take $\hat{m}$ to be $\hat{F}$-invariant. Then, for $\hat{\mu}$-almost every $\hat{q} \in \hat{M}$, the sequence $(A^n(q_n)_* m_{q_n})_n$ converges to $\hat{m}_{\hat{q}}$ in the weak* topology.*

Proof. Let $\phi : \mathrm{Grass}(l, d) \to \mathbb{R}$ be any continuous function. Define

$$I_n : M \to \mathbb{R}, \quad I_n(q) = \int \phi \circ A^n(q)\, dm_q$$

$$\hat{I}_n : \hat{M} \to \mathbb{R}, \quad \hat{I}_n(\hat{q}) = \int \phi \circ A^n(q_n)\, dm_{q_n} = I_n(q_n).$$

Thus $\hat{I}_n = I_n \circ (P \times \mathrm{id}_K) \circ \hat{f}^{-n}$. Let $\mathfrak{C}$ be the Borel σ-algebra of M and, for each $n \geq 0$, consider the σ-algebra $\mathfrak{C}_n = \hat{f}^n((P \times \mathrm{id}_K)^{-1}(\mathfrak{C}))$ of subsets of $\hat{M}$. Note that the union $\cup_{n \in \mathbb{N}} \mathfrak{C}_n$ generates the Borel σ-algebra of $\hat{M}$.

For each $k \geq 1$, let $\{\mu_q^k : q \in M\}$ be the disintegration of μ with respect to the partition $\{f^{-k}(q) : q \in M\}$ of M into pre-images under f^k. Since f^k is countable-to-one, we may express the disintegration explicitly in terms of the Jacobian:

$$\mu_q^k = \sum_{y \in f^{-k}(q)} \frac{1}{J_\mu f^k(y)} \delta_y \text{ for } \mu\text{-almost every } q \in M. \tag{18}$$

Lemma 4.4. *$I_n(q) = \int I_{n+k}(z)\, d\mu_q^k(z)$ for μ-almost every $q \in M$ and every $n \geq 0$.*

Proof. Since m is F-invariant, we may use Proposition 4.1 to get that

$$\begin{aligned} I_n(q) &= \int \phi \circ A^n(q)\, d\Big(\int A^k(z)_* m_z\, d\mu_q^k(z) \Big) \\ &= \int \int \phi \circ A^n(q)\, d\big(A^k(z)_* m_z\, d\mu_q^k(z) \big) \\ &= \int \int \phi \circ A^{n+k}(z)\, dm_z\, d\mu_q^k(z) = \int I_{n+k}(z)\, d\mu_q^k(z), \end{aligned}$$

as claimed. □

Lemma 4.5. *The sequence $(\hat{I}_n, \mathfrak{C}_n)_n$ is a martingale.*

Proof. It is clear from the definition that every I_n is $\mathfrak{C}_0$-measurable. Hence, $\hat{I}_n$ is $\mathfrak{C}_n$-measurable for every $n \geq 0$. We are left to check that

$$\int \hat{I}_n(\hat{q}) \hat{\psi}(\hat{q})\, d\hat{\mu}(\hat{q}) = \int \hat{I}_{n+k}(\hat{q}) \psi(\hat{q})\, d\hat{\mu}(\hat{q}) \tag{19}$$

for any $\mathfrak{C}_n$-measurable function $\hat\psi : \hat M \to \mathbb{R}$ and any $n \ge 0$ and $k \ge 1$.

By definition, we may write $\hat\psi = \psi_n \circ (P \times \mathrm{id}_K) \circ \hat f^{-n}$ for some $\mathfrak{C}_0$-measurable function $\hat\psi_n : M \to \mathbb{R}$. Since $\hat\mu$ is $\hat f$-invariant and μ is its projection,

$$\int \hat I_n(\hat q)\hat\psi(\hat q)\, d\hat\mu(\hat q) = \int I_n(q)\psi_n(q)\, d\mu(q) \text{ and}$$
$$\int \hat I_{n+k}(\hat q)\psi(\hat q)\, d\hat\mu(\hat q) = \int I_{n+k}(q)\psi_n(f^k(q))\, d\mu(q)$$

(note that $\hat\psi = \psi_n \circ f^k \circ (P \times \mathrm{id}_K) \circ \hat f^{-(n+k)}$). By Lemma 4.4 and the definition of μ_q^k,

$$\begin{aligned}\int I_n(q)\psi_n(q)\, d\mu(q) &= \int\int I_{n+k}(z)\, d\mu_q^k(z)\psi_n(q)\, d\mu(q)\\ &= \int\int I_{n+k}(z)\psi_n(f^k(z))\, d\mu_q^k(z)\, d\mu(q)\\ &= \int I_{n+k}(q)\psi_n(f^k(q))\, d\mu(q).\end{aligned}$$

Combining these identities, one gets the property (19). □

So, we may use the martingale convergence theorem to conclude that, given any continuous function $\phi : \mathrm{Grass}(l,d) \to \mathbb{R}$,

$$\hat I_\phi(\hat q) = \lim_{n\to\infty} \hat I_n(\hat q)$$

exists for $\hat\mu$-almost every $\hat q \in \hat M$. So, using the fact that the space $C(\mathrm{Grass}(l,d),\mathbb{R})$ of continuous functions is separable, one may find a set $\hat M_0 \subset \hat M$ with full $\hat\mu$-measure such that $\hat I_\phi(\hat q)$ is defined for every $\hat q \in \hat M_0$ and $\phi \in C(\mathrm{Grass}(l,d),\mathbb{R})$. Then, by Riesz-Markov, for each $\hat q \in \hat M_0$ there exists a probability measure $\tilde m_{\hat q}$ on $\mathrm{Grass}(l,d)$ such that

$$\int \phi\, d\tilde m_{\hat q} = \hat I_\phi(\hat q) = \lim_{n\to\infty} \hat I_n(\hat q) = \lim_{n\to\infty} \int \phi \circ A^n(q_n)\, dm_{q_n}(v)$$

for any $g \in C(\mathrm{Grass}(l,d),\mathbb{R})$. In other words, $(A^n(q_n)_* m_{q_n})_n$ converges to $\tilde m_{\hat q}$ in the weak* topology.

We are left to prove that $\tilde m_{\hat q} = \hat m_{\hat q}$ for $\hat\mu$-almost every $\hat q$. Let $n \ge 0$ and $\hat\psi : \hat M \to \mathbb{R}$ be any $\mathfrak{C}_n$-measurable function. Taking the limit as $k \to \infty$ in (19), we get that

$$\int \hat I_\phi(\hat q)\hat\psi(\hat q)\, d\hat\mu(\hat q) = \int \hat I_n(\hat q)\hat\psi(\hat q)\, d\hat\mu(\hat q)$$

for every $\phi \in C(\mathrm{Grass}(l,d),\mathbb{R})$. In other words,

$$\int\int \hat\psi(\hat q)\phi(\xi)\, d\tilde m_{\hat q}(\xi)\, d\hat\mu(\hat q) = \int\int \hat\psi(\hat q)\phi(A^n(q_n)\xi)\, dm_{q_n}(\xi)\, d\hat\mu(\hat q).$$

Since $\hat\mu$ is $\hat f$-invariant and $\hat A(\hat q) = A(q_0)$ for every $\hat q \in \hat M$, the right-hand side of the previous identity may be rewritten as

$$\begin{aligned}&\int\int \hat\psi(\hat f^n(\hat q))\phi(A^n(q_0)\xi)\, dm_{q_0}(\xi)\, d\hat\mu(\hat q)\\ &\qquad = \int\int \hat\psi(\hat f^n(\hat q))\phi(\hat A^n(\hat q)\xi)\, d\hat m_{\hat q}(\xi)\, d\hat\mu(\hat q).\end{aligned}$$

Finally, this last expression may be rewritten as

$$\int\int \hat\psi(\hat f^n(\hat q))\phi(\hat A^n(\hat q)\xi)\, d\hat m_{\hat q}(\xi)d\hat\mu(\hat q) = \int\int \hat\psi(\hat q)\phi(\xi)\, d\hat m_{\hat q}(\xi)d\hat\mu(\hat q),$$

because $\hat m$ is $\hat F$-invariant. The last three identities prove that

$$\int\int \hat\psi(\hat q)\phi(\xi)\, d\tilde m_{\hat q}(\xi)\, d\hat\mu(\hat q) = \int\int \hat\psi(\hat q)\phi(\xi)\, d\hat m_{\hat q}(\xi)d\hat\mu(\hat q). \tag{20}$$

This relation extends to any function on $\hat M \times \mathrm{Grass}(l,d)$ which is a linear combination of functions $\hat\psi \times \phi$ such that $\hat\psi$ is $\mathfrak{C}_n$-measurable for some $n \geq 1$ and ϕ is continuous. Since these linear combinations are dense in the set of all bounded measurable functions on $\hat M \times \mathrm{Grass}(l,d)$, it follows that (20) holds for every bounded measurable functions on $\hat M \times \mathrm{Grass}(l,d)$. This means that $\{\tilde m_{\hat q} : \hat q \in \hat M\}$ is a disintegration of $\hat m$. So, by essential uniqueness, $\tilde m_{\hat q} = \hat m_{\hat q}$ for $\hat\mu$-almost every $\hat q$. □

Proposition 4.6. *For $\hat\mu$-almost every $\hat q \in \hat M$, any $k \geq 1$ and any choice of points $y_{n,k}$ such that $f^k(y_{n,k}) = q_n$ and $\{y_{n,k} : n \geq 0\}$ is contained in some k-cylinder,*

$$\lim_{n\to\infty} A^n(q_n)_* m_{q_n} = \lim_{n\to\infty} A^{n+k}(y_{n,k})_* m_{y_{n,k}}.$$

Proof. For every $n \geq 0$ and $k \geq 1$, define

$$S_{n,k} = \int\int \big(I_{n+k}(z) - \hat I_n(\hat q)\big)^2\, d\mu^k_{q_n}(y)\, d\hat\mu(\hat q).$$

Lemma 4.7. *For every $n \geq 0$ and $k \geq 1$,*

$$S_{n,k} = \int \hat I_{n+k}(\hat q)^2\, d\hat\mu(\hat q) - \int \hat I_n(\hat q)^2\, d\hat\mu(\hat q).$$

Proof. Using the definitions of μ^k_q and $\hat I_{n+k}$, together with the fact that $\hat\mu$ is $\hat f$-invariant,

$$\begin{aligned}\int\int I_{n+k}(z)^2\, d\mu^k_{q_n}(z)\, d\hat\mu(\hat q) &= \int I_{n+k}(q_n)^2\, d\hat\mu(\hat q)\\ &= \int \hat I_{n+k}(\hat f^k(\hat q))^2\, d\hat\mu(\hat q) = \int \hat I_{n+k}(\hat q)^2\, d\hat\mu(\hat q).\end{aligned}$$

On the other hand, using Lemma 4.4,

$$\begin{aligned}\int \hat I_n(\hat q)\int I_{n+k}(z)\, d\mu^k_{q_n}(z)\, d\hat\mu(\hat q) &= \int \hat I_n(\hat q) I_n(q_n)\, d\hat\mu(\hat q)\\ &= \int \hat I_n(\hat q)^2\, d\hat\mu(\hat q).\end{aligned}$$

The claim follows. □

It follows from Lemma 4.7 that, for any $s \geq 0$,

$$\begin{aligned}\sum_{n=0}^{s}\int\int \big(I_{n+k}(y) - \hat I_n(\hat q)\big)^2\, d\mu^k_{q_n}(y)\, d\hat\mu(\hat q)&\\ = \sum_{j=0}^{k-1}\int \hat I_{s-j}(\hat q)^2\, d\hat\mu(\hat q) - \int \hat I_j(\hat q)^2\, d\hat\mu(\hat q).&\end{aligned}$$

The right-hand side is uniformly bounded (by $2k \sup |g|^2$). It follows that

$$\lim_{n\to\infty} \int \big(I_{n+k}(y) - \hat{I}_n(\hat{q})\big)^2 \, d\mu^k_{q_n}(y) = 0 \text{ for } \hat{\mu}\text{-almost every } \hat{q} \in \hat{M}.$$

In view of (18), this implies that

$$\lim_{n\to\infty} \frac{1}{J_\mu f^k(y_{n,k})} \big(I_{n+k}(y_{n,k}) - \hat{I}_n(\hat{q})\big)^2 = 0 \text{ for } \hat{\mu}\text{-almost every } \hat{q} \in \hat{M}.$$

Finally, since the Jacobian is bounded from above on every k-cylinder (Lemma 3.8), we deduce that

$$\Big(\int \phi A^{n+k}(y_{n,k}) \, dm_{y_{n,k}} - \int \phi A^n(q_n) \, dm_{q_n} \Big)^2 = \big(I_{n+k}(y_{n,k}) - \hat{I}_n(\hat{q})\big)^2$$

converges to 0 as $n \to \infty$, for $\hat{\mu}$-almost every $\hat{q} \in \hat{M}$.

Then, using the fact that $C(\mathrm{Grass}(l,d), \mathbb{R})$ is separable, one can find a full $\hat{\mu}$-measure set of values of $\hat{q}$ such that the latter convergence holds for every $\phi \in C(\mathrm{Grass}(l,d), \mathbb{R})$. □

The argument extends immediately to the case when $\{y_{n,k} : n \geq 0\}$ is contained in any finite union of k-cylinders.

5. Properties of u-states

A probability measure $\hat{m}$ on $\hat{M} \times \mathrm{Grass}(l,d)$ is called a *u-state* of $\hat{F}$ if there exist a disintegration $\{\hat{m}_{\hat{q}} : \hat{q} \in \hat{M}\}$ with respect to the partition $\big\{\{\hat{q}\} \times \mathrm{Grass}(l,d) : \hat{q} \in \hat{M}\big\}$ which is invariant under unstable holonomy:

$$\hat{m}_{\hat{q}} = H^u_{\hat{b},\hat{q}_*} \hat{m}_{\hat{b}} \text{ for every } \hat{b}, \hat{q} \in \tilde{M} \text{ with } \hat{q} \in W^{uu}_{loc}(\hat{b}), \tag{21}$$

where $\tilde{M} \subset \hat{M}$ is some full measure set. Let $\pi : \hat{M} \times \mathrm{Grass}(l,d) \to \hat{M}$ be the canonical projection.

Proposition 5.1. *There is some $\hat{F}$-invariant u-state $\hat{m}$ that projects down to $\hat{\mu}$ under π.*

This is entirely analogous to Proposition 4.2 of [3], and so we only outline the proof. The idea is to fix some $\hat{x} \in \hat{\Sigma}$ and define a homeomorphism between the measures in $\{\hat{x}\} \times W^s_{loc}(\hat{x}) \times K$ that projects to μ^s and the u-states. Using that this space in compact we have that the space of u-states measures is also compact and $\hat{F}_*$-invariant, so we have that any accumulation point of $n^{-1} \sum_{j=0}^{n-1} \hat{F}^j_* \hat{m}$ is also a u-state.

In the remainder of this section, $\hat{m}$ denotes any $\hat{F}$-invariant u-state that projects down to $\hat{\mu}$ under π, and $\{\hat{m}_{\hat{q}} : \hat{q} \in \hat{M}\}$ is taken to be a disintegration as in (21).

5.1. **Bounded distortion.** Let $\pi_1 : \hat{M} \to \hat{\Sigma}$ be the canonical projection $\pi_1(\hat{x}, t) = \hat{x}$ and denote $\hat{\nu} = \pi_{1*}\hat{\mu}$. Equivalently,

$$\hat{\nu}(E) = \int_{E \times K} \hat{\rho}(x^s, x, t) \, d\mu^s(x^s) \, d\mu^u(x) \, d\mu^c(t)$$

for any measurable set $E \subset \hat{\Sigma}$. For each $x \in \Sigma$, define $\hat{\nu}_x$ to be the normalization of

$$\mu^s \int \hat{\rho}(\cdot, x, t) \, d\mu^c(t).$$

Then $\{\hat{\nu}_x : x \in \Sigma\}$ is a continuous disintegration of $\hat{\nu}$ with respect to the partition into local stable sets $W^s_{loc}(\hat{x})$.

The measure $\hat{\nu}$ satisfies the properties of local product structure, boundedness and continuity in [3, Section 1.2]. In what follows, we recall a few results about this type of measures that we will use later.

For each $x^u \in \Sigma^+$ and $k \geq 1$ let the *backward average* measure μ^u_{k,x^u} of the map σ be defined by

$$\mu^u_{k,x^u} = \sum_{\sigma^k(z)=x^u} \frac{1}{J\sigma^k(z)} \delta_z$$

where $J\sigma^k : \Sigma^+ \to \mathbb{R}$ is the Jacobian of μ^u with respect to σ^k.

Lemma 5.2. *For any cylinder $I^u = [0; \iota_0, \ldots, \iota_{k-1}] \subset \Sigma^+$ and $z^u \in I^u$,*

$$\hat{\sigma}^k_* \hat{\nu}_{z^u} = J\sigma^k(z^u)(\hat{\nu}_{\sigma^k(z^u)} \mid I^s)$$

where $\{\hat{\nu}_{z^u} : z^u \in \Sigma^+\}$ is the disintegration of $\hat{\nu}$ with respect to the partition $\{\Sigma^- \times \{z^u\} : z^u \in \Sigma^+\}$.

Proof. Analogous to [3, Lemma 2.6]. □

Lemma 5.3. *For every cylinder $[J] \subset \Sigma^+$ and $x^u \in \Sigma^+$,*

$$\begin{aligned} \kappa\mu^u([J]) &\geq \limsup_n \frac{1}{n} \sum_{k=0}^{n-1} \mu^u_{k,x^u}([J]) \\ &\geq \liminf_n \frac{1}{n} \sum_{k=0}^{n-1} \mu^u_{k,x^u}([J]) \geq \frac{1}{\kappa}\mu^u([J]). \end{aligned}$$

Proof. Analogous to [3, Lemma 2.7]. □

As a direct consequence, for every cylinder $[J] \subset \Sigma^+$ and $x^u \in \Sigma^+$,

$$\limsup_k \mu^u_{k,x^u}([J]) \geq \frac{1}{\kappa}\mu^u([J]). \tag{22}$$

5.2. **Estimating the Jacobians.** For every $x \in \Sigma$ let

$$F_x : K \times \mathrm{Grass}(l,d) \to K \times \mathrm{Grass}(l,d), \quad F_x(t,V) = (f_x(t), A(x,t)V)$$

and for every $\hat{x}, \hat{y} \in \hat{\Sigma}$ in the same unstable set let

$$H_{\hat{x},\hat{y}} : K \times \mathrm{Grass}(l,d) \to K \times \mathrm{Grass}(l,d)$$

be defined by

$$H_{\hat{x},\hat{y}}(t,V) = (h^u_{\hat{x},\hat{y}}(t), H^u_{(\hat{x},t)(\hat{y},h^u_{\hat{x},\hat{y}}(t))}V).$$

Observe that $F_{\hat{y}} \circ H_{\hat{x},\hat{y}} = H_{\hat{\sigma}(\hat{x}),\hat{\sigma}(\hat{y})} \circ F_{\hat{x}}$. Now define $\{\hat{m}_{\hat{x}} : \hat{x} \in \hat{\Sigma}\}$ as

$$\hat{m}_{\hat{x}} = \int \hat{m}_{\hat{x},t} \, d\hat{\mu}^c_{\hat{x}}(t).$$

Observe that for any $\varphi : K \times \mathrm{Grass}(l,d) \mapsto \mathbb{R}$,

$$\begin{aligned}\int \varphi d\big(H_{\hat{x},\hat{y}*}\hat{m}_{\hat{x}}\big) &= \int \varphi\big(h^u_{\hat{x},\hat{y}}(t), H^u_{(\hat{x},t)(\hat{y},h^u_{\hat{x},\hat{y}}(t))}V\big)d\,\hat{m}_{\hat{x},t}(V)d\,\hat{\mu}^c_{\hat{x}}(t)\\ &= \int \varphi\big(h^u_{\hat{x},\hat{y}}(t), V\big)d\,\big(H^u_{(\hat{x},t)(\hat{y},h^u_{\hat{x},\hat{y}}(t))*}\hat{m}_{\hat{x},t}\big)(V)d\,\hat{\mu}^c_{\hat{x}}(t)\\ &= \int \varphi\big(h^u_{\hat{x},\hat{y}}(t), V\big)d\,\hat{m}_{\hat{y},h^u_{\hat{x},\hat{y}}(t)}(V)d\,\hat{\mu}^c_{\hat{x}}(t)\\ &= \int \varphi\big(t, V\big)d\,\hat{m}_{\hat{y},t}(V)d\,(h^u_{\hat{x},\hat{y}*}\hat{\mu}^c_{\hat{x}})(t)\\ &= \int \varphi\big(t, V\big)d\,\hat{m}_{\hat{y},t}(V)d\,\hat{\mu}^c_{\hat{y}}(t),\end{aligned}$$

because $\hat{m}$ is a u-state and $\{\hat{\mu}^c_{\hat{x}} : \hat{x} \in \hat{\Sigma}\}$ is u-invariant. It is also easy to see that $\{\hat{m}_{\hat{x}} : \hat{x} \in \hat{\Sigma}\}$ is a disintegration of $\hat{m}$ with respect to the partition $\{\hat{x} \times K \times \mathrm{Grass}(l,d) : \hat{x} \in \hat{\Sigma}\}$.

The main point with the next corollary is that the conclusion is stated for *every* $x \in \Sigma$.

Corollary 5.4. *If $\{\hat{m}_{\hat{x}} : \hat{x} \in \hat{\Sigma}\}$ is a disintegration of an invariant u-state $\hat{m}$ then*

$$\hat{m}_{\hat{\sigma}^n(\hat{x})} = F^n_{x\,*}\hat{m}_{\hat{x}}$$

for every $n \geq 1$, every $x \in \Sigma$, and $\hat{\nu}_x$-almost every $\hat{x} \in W^s_{loc}(x)$.

Proof. Since $\hat{m}$ is $\hat{F}$-invariant, the equality is true for all $n \geq 1$ and $\hat{\nu}$-almost all $\hat{z} \in \hat{\Sigma}$ or, equivalently, for $\hat{\nu}_z$-almost every $\hat{z} \in W^s_{loc}(z)$ and ν-almost every $z \in \Sigma$. Consider an arbitrary point $x \in \Sigma$. Since ν is positive on open sets, x may be approximated by points z such that

$$\hat{m}_{\hat{\sigma}^n(\hat{z})} = F^n_{z\,*}\hat{m}_{\hat{z}}$$

for every $n \geq 1$ and $\hat{\mu}_z$-almost every $\hat{z} \in W^s_{loc}(z)$. Since the conditional probabilities of $\hat{m}$ are invariant under unstable holonomies, it follows that

$$\hat{m}_{\hat{\sigma}^n(\hat{x})} = (H_{\hat{\sigma}^n(z),\hat{\sigma}^n(x)})_* F^n_{z\,*}\hat{m}_{\hat{z}} = F^n_{x\,*}(H_{\hat{z},\hat{x}})_*\hat{m}_{\hat{z}} = F^n_{x\,*}\hat{m}_{\hat{x}}$$

for $\hat{\mu}_z$-almost every $\hat{z} \in W^s_{loc}(z)$, where $\hat{x}$ is the unique point in $W^s_{loc}(x) \cap W^u_{loc}(\hat{z})$. Since the measures $\hat{\mu}_x$ and $\hat{\mu}_z$ are equivalent, this is the same as saying that the last equality holds for $\hat{\mu}_x$-almost every $\hat{x} \in W^s_{loc}(x)$, as claimed. □

5.3. **L^1-continuity of conditional probabilities.** Recall that

$$\mu = (P \times \mathrm{id}_K)_*\,\hat{\mu} \quad \text{and} \quad m = \big(P \times \mathrm{id}_K \times \mathrm{id}_{\mathrm{Grass}(l,d)}\big)_*\,\hat{m}.$$

Let $\{m_x : x \in \Sigma\}$ and $\{m_{x,t} : (x,t) \in M\}$ be disintegrations of m with respect to the partitions $\{\{x\} \times K \times \mathrm{Grass}(l,d), x \in \Sigma\}$ and $\{\{(x,t)\} \times \mathrm{Grass}(l,d), (x,t) \in \Sigma \times K\}$, respectively. Thus each m_x is a probability measure on $K \times \mathrm{Grass}(l,d)$ and each $m_{x,t}$ is a probability measure on $\mathrm{Grass}(l,d)$.

It is easy to check that $x \mapsto m_x$ may be chosen to be continuous with respect to the weak* topology, indeed, we will do that in a while. The corresponding statement for $x \mapsto m_{x,t}$ is false, in general. However, the main point of this section is to show that the family $\{m_{x,t} : (x,t) \in M\}$ does have some continuity property:

Proposition 5.5. *Let $(x_n)_n$ be a sequence in Σ converging to some $x \in \Sigma$. Then there exists a sub-sequence $(x_{n_k})_k$ such that*

$$m_{x_{n_k},t} \to m_{x,t} \text{ as } k \to \infty$$

in the weak topology, for μ^c-almost every $t \in K$.*

We will deduce this from a somewhat stronger L^1-continuity result, whose precise statement will be given in Proposition 5.9. The key ingredient in the proofs is a result about maps on geodesically convex metric spaces that we are going to state in Lemma 5.7 and which will also be useful at latter stages of our arguments.

Definition 5.6. A metric space N is *geodesically convex* if there exists $\tau \geq 1$ such that for every $u, v \in N$ there exist a continuous path $\lambda : [0,1] \to N$ with $\lambda(0) = u$, $\lambda(1) = v$ and

$$\operatorname{dist}_N(\lambda(t), \lambda(s)) \leq \tau \operatorname{dist}_N(u, v) \text{ for every } s, t \in [0,1].$$

Geodesically convex metric spaces include convex subsets of a Banach space, path connected compact metric spaces and complete connected Riemannian manifolds, among other examples. The spaces of maps with values in a geodesically convex metric space are analyzed in Appendix A.

Lemma 5.7. *Let L be a geodesically convex metric space and take $(K, \mathfrak{B}_K, \mu_K)$ to be a probability space such that K is a normal topological space, $\mathfrak{B}_K$ is the Borel σ-algebra of K and μ_K is a regular measure.*

Let $H_{j,t} : L \to L$ and $h_j : K \to K$, with $j \in \mathbb{N}$ and $t \in K$, be such that

$$\big(H_{j,t}(x)\big)_j \to x \quad \text{and} \quad \big(h_j(t)\big)_j \to t,$$

uniformly in $t \in K$ and $x \in L$ and, moreover, the Jacobian $Jh_j(t)$ of each h_j with respect to μ_K is uniformly bounded. Then

$$\lim_j \int \operatorname{dist}_L\big(\psi(t), H_{j,t} \circ \psi \circ h_j(t)\big)\, d\mu_K(t) = 0$$

for every bounded measurable map $\psi : K \to L$.

Proof. Take $j \in \mathbb{N}$ to be sufficiently large that $d_L(H_{j,t}(x), x) < \epsilon/4$ for every t and x. Then,

$$\begin{aligned}\int \operatorname{dist}_L(\psi, H_{j,t} \circ \psi \circ h_j)\, d\mu_K &\leq \int \big(\operatorname{dist}_L(\psi, \psi \circ h_j) \\ &\qquad + \operatorname{dist}_L(\psi \circ h_j, H_{j,t} \circ \psi \circ h_j)\big)\, d\mu_K \\ &\leq \int \operatorname{dist}_L(\psi, \psi \circ h_j)\, d\mu_K + \frac{\epsilon}{4}.\end{aligned}$$

Let $C > 1$ be a uniform bound for $Jh_j(t)$. By Proposition A.1, given $\epsilon > 0$ there exists a continuous map $\tilde{\psi} : K \to L$ such that

$$\int \operatorname{dist}_L(\tilde{\psi}, \psi)\, d\mu_K < \frac{\epsilon}{4C}.$$

Then, by change of variables,

$$\int \operatorname{dist}_L(\tilde{\psi} \circ h_j, \psi \circ h_j)\, d\mu_K \leq C \int \operatorname{dist}_L(\tilde{\psi}, \psi)\, d\mu_K < \frac{\epsilon}{4}.$$

Then

$$\begin{aligned}\int &\operatorname{dist}_L(\psi, \psi \circ h_j)\, d\mu_K \\ &\le \int \big(\operatorname{dist}_L(\psi, \tilde\psi) + \operatorname{dist}_L(\tilde\psi, \tilde\psi \circ h_j) + \operatorname{dist}_L(\tilde\psi \circ h_j, \psi \circ h_j) \big)\, d\mu_K \\ &\le \int \operatorname{dist}_L(\tilde\psi, \tilde\psi \circ h_j)\, d\mu_K + \frac{\epsilon}{2}.\end{aligned}$$

By the continuity of $\tilde\psi$, increasing j if necessary,

$$\operatorname{dist}_L(\tilde\psi(t), \tilde\psi \circ h_j(t)) < \frac{\epsilon}{4} \text{ for every } t \in K.$$

The conclusion follows from these inequalities. □

Lemma 5.8. *Let $(x_n)_n$ be a sequence in Σ converging to some $x \in \Sigma$ and let $(j_n)_n$ be a sequence of integer numbers going to infinity. Assume that $(f^{j_n}_{x_n})_n$ converges uniformly to some $g : K \to K$ and $(\sigma^{j_n}(x_n))_n$ converges to some $z \in \Sigma$. Then g is absolutely continuous with respect to μ^c and it has bounded Jacobian.*

Proof. By Lemma 3.5 we have that

$$(f^{j_n}_{x_n})_* \mu^c_{x_n} = \mu^c_{\sigma^{j_n}(x_n)}.$$

Taking $n \to \infty$ we get that

$$g_* \mu^c_x = \mu^c_z,$$

which implies that $Jg_{\mu^c} = \frac{\varrho(z,t)}{\varrho(x,t)}$ is uniformly bounded. □

Proposition 5.9. *Let $\varphi : \operatorname{Grass}(l,d) \to \mathbb{R}$ be a continuous function, $(x_n)_n$ be a sequence in Σ converging to some $x \in \Sigma$ and $(j_n)_n$ be a sequence of integer numbers such that $(f^{j_n}_{x_n})_n$ converges uniformly to some $g : K \to K$. Then $\int \varphi dm_{x_n, f^{j_n}_{x_n}(t)}$ converges to $\int \varphi\, dm_{x,g(t)}$ in $L^1(\mu^c)$.*

Proof. Denote $t_n = f^{j_n}_{x_n}(t)$. Fix $x^s \in \Sigma^-$ and let

$$h^u_{n,x^s}(t) = h^u_{(x^s,x_n),(x^s,x)}(t_n)$$

and

$$H^u_{(x^s,x,h^u_{n,x^s}(t)),(x^s,x_n,t_n)} = H^u_{n,x^s,t}.$$

Note that

$$\hat m_{x^s,x_n,t_n} = \big(H^u_{n,x^s,t}\big)_*\, \hat m_{x^s,x,h^u_{n,y}(t_n)}.$$

So, applying Lemma 5.7 with $L = \mathcal{M}$ (the compact metric space of probability measures on $\operatorname{Grass}(l,d)$), $H_{j,t} = \big(H^u_{j,x^s,t}\big)_*$, $h_j = h^u_{j,x^s}(t_j)$ and

$$\psi : K \to \mathcal{M}, \quad \psi(t) = \hat m_{x^s,x,g(t)},$$

we get that the sequence $\int \varphi d\hat m_{x^s,x_n,f^{j_n}_{x_n}(t)}$ converges to $\int \varphi d\hat m_{y,x,g(t)}$ in the space $L^1(\mu^c)$.

By the definition of disintegration,

$$m_{x,t} = \int \hat\rho(x^s,x,t) \hat m_{x^s,x,t}\, d\mu^s(x^s)$$

and so

$$\int |\int \varphi(v) dm_{x_n,t_n} - \int \varphi(v) dm_{x,g(t)}|\, d\mu^c \leq \int\int |\int \varphi\rho(x^s, x_n, t_n) d\hat{m}_{x^s,x_n,t_n} - \int \varphi\rho(x^s, x, g(t)) d\hat{m}_{x^s,x,g(t)}|\, d\mu^c\, d\mu^s.$$

So, noting that the integrand goes to zero as $n \to \infty$, for every $x^s \in \Sigma^-$, the dominated convergence theorem ensures that

$$\lim_{n\to\infty} \int |\int \varphi(v)\, dm_{x_n,t_n} - \int \varphi(v)\, dm_{x,g(t)}|\, d\mu^c = 0,$$

as we wanted to prove. □

This proposition implies that, given any $(x_n)_n \to x$ and any continuous $\varphi : \mathrm{Grass}(l,d) \to \mathbb{R}$, there exist a sub-sequence $(n_k)_k$ such that

$$\int \varphi\, dm_{x_{n_k},t} \to \int \varphi\, dm_{x,t} \text{ for } \mu^c - \text{almost every } t.$$

Moreover, since the space of continuous functions is separable, one can use a diagonal argument (see e.g. the proof of [21, Proposition 2.1.6]) to construct such a sub-sequence independent of φ. In other words,

$$m_{x_{n_k},t} \to m_{x,t} \text{ in the weak}^*\text{-topology, for } \mu^c - \text{almost every } t.$$

This proves Proposition 5.5.

Corollary 5.10. *The disintegration $\{m_x : x \in \Sigma\}$ is continuous.*

Proof. Let $\varphi : K \times \mathrm{Grass}(l,d) \to \mathbb{R}$ be a continuous function. Given any $(x_n)_n \to x$, we have that

$$\begin{aligned} &|\int \varphi dm_{x_n} - \int \varphi dm_x| \\ &= |\int\int \varphi(t,v)\, dm_{x_n,t}(v)\rho(x_n,t)\, d\mu^c(t) \\ &\qquad - \int\int \varphi(t,v)\, dm_{x,t}(v)\rho(x,t)\, d\mu^c(t)| \\ &\leq \int |\int \varphi(t,v)\rho(x_n,t)\, dm_{x_n,t}(v) - \int \varphi(t,v)\rho(x,t)\, dm_{x,t}(v)|\, d\mu^c(t). \end{aligned}$$

By Proposition 5.5, up to restricting to a subsequence, we may suppose that $(m_{x_n,t})_n$ converges to $m_{x,t}$ in the weak* sense, for μ^c-almost every t. Then

$$\int \varphi(t,v)\rho(x_n,t)\, dm_{x_n,t}(v) \to \int \varphi(t,v)\rho(x,t)\, dm_{x,t}(v)$$

for μ^c-almost every t. To get the conclusion it suffices to use this observation in the previous inequality, together with dominated convergence. □

Corollary 5.11. *We have $m_x = \int (F_y^k)_* m_y\, d\mu_x^k(y)$ for every $x \in \Sigma$ and $k \geq 1$.*

Proof. The F-invariance of m gives that $m_x = \int (F_y^k)_* m_y\, d\mu_x^k(y)$ for ν-almost every x, the continuity of the disintegration implies that this extends to every $x \in \Sigma$. □

6. Dual graphs of Grassmanian sections

Let $w_1, \dots, w_l$ be a basis of a given subspace $W \in \mathrm{Grass}(l, d)$. The exterior product $w_1 \wedge \cdots \wedge w_l$ depends on the choice of the basis, but its projective class does not. Thus we have a well-defined map

$$\mathrm{Grass}(l, d) \hookrightarrow \mathbb{P}\Lambda^l\mathbb{C}^d, \quad W \mapsto [w_1 \wedge \cdots \wedge w_l], \tag{23}$$

which can be checked to be an embedding: it is called the *Plücker embedding* of $\mathrm{Grass}(l, d)$. The image is the projectivization of the *space of l-vectors*

$$\Lambda^l_v(\mathbb{C}^d) = \{w_1 \wedge w_2 \wedge \cdots \wedge w_l \in \Lambda^l\mathbb{C}^d : w_i \in \mathbb{C}^d \text{ for } 1 \le i \le l\},$$

which we denote by $\mathbb{P}\Lambda^l_v(\mathbb{C}^d)$. This is a closed subset of $\mathbb{P}\Lambda^l\mathbb{C}^d$, and it is invariant under the action induced on $\mathbb{P}\Lambda^l\mathbb{C}^d$ by any linear linear transformation $B : \mathbb{C}^d \to \mathbb{C}^d$. See [3, Section 2] for more information about l-vectors.

The *geometric hyperplane* $\mathfrak{H}V \subset \mathrm{Grass}(l, d)$ associated with a subspace $V \in \mathrm{Grass}(d - l, d)$ is the set $\mathfrak{H}V$ of all subspaces $W \in \mathrm{Grass}(l, d)$ which are *not* in general position relative to V. In other words,

$$\mathfrak{H}V = \{W \in \mathrm{Grass}(l, d) : W \cap V \neq \{0\}\}.$$

This may also be formulated using the Plücker embedding (23): if v is any $(d-l)$-vector representing V, then $\mathfrak{H}V$ consists of the subspaces $W \in \mathrm{Grass}(l, d)$ represented by l-vectors w such $v \wedge w \neq 0$.

In what follows, let $\mathfrak{H}\,\mathrm{Sec}$ denote the space of measurable sections $\mathcal{V} : K \to \mathrm{Grass}(d - l, d)$. We define the *dual graph* of each $\mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}$ to be

$$\mathrm{graph}\,\mathfrak{H}\mathcal{V} = \{(t, v) \in K \times \mathrm{Grass}(l, d) : v \in \mathfrak{H}\mathcal{V}(t)\}.$$

Let $\hat{m}$ be any u-state on $\hat{M} \times \mathrm{Grass}(l, d)$, m be its projection to $M \times \mathrm{Grass}(l, d)$ and $\{m_x : x \in \Sigma\}$ be the Rokhlin disintegration of m along the fibers $K \times \mathrm{Grass}(l, d)$ (recall Section 5.3). The purpose of this section is to prove the following fact:

Proposition 6.1. *We have $m_x(\mathrm{graph}\,\mathfrak{H}\mathcal{V}) = 0$ for any u-state $\hat{m}$, any $\mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}$ and every $x \in \Sigma$.*

6.1. **Graphs have measure zero.** Starting the proof of Proposition 6.1, recall that each m_x is a probability measure on $K \times \mathrm{Grass}(l, d)$, and

$$m_x = \int m_{x,t}\varrho(x, t)\, d\mu^c(t),$$

where each $m_{x,t}$ is a probability measure on $\mathrm{Grass}(l, d)$. Recall also that $x \mapsto m_x$ is continuous, by Corollary 5.10.

Let $x \in \Sigma$ be fixed for the time being, and consider the functions

$$G : K \times \mathrm{Grass}(d - l, d) \mapsto \mathbb{R}, \quad G\,(t, V) = m_{x,t}(\mathfrak{H}V) \text{ and}$$
$$g : K \to \mathbb{R}, \quad g(t) = \sup\{m_{x,t}(\mathfrak{H}Z) : Z \in \mathrm{Grass}(d - l, d)\}.$$

Lemma 6.2. *$G : K \times \mathrm{Grass}(d - l, d) \mapsto \mathbb{R}$ and $g(t) : K \mapsto \mathbb{R}$ are measurable functions.*

Proof. Let $\mathcal{P}^1 \prec \mathcal{P}^2 \prec \cdots$ be an increasing sequence of finite partitions of $\mathrm{Grass}(d-l, d)$ such that $\mathcal{P} = \vee_{i\in\mathbb{N}}\mathcal{P}^i$ is the partition into points (that such a sequence exists

is clear, e.g., because the Grassmannian is compact). Write $\mathcal{P}^i = \{P_1^i, \cdots, P_{n_i}^i\}$ and then define

$$G_n : K \times \mathrm{Grass}(d-l,d) \mapsto \mathbb{R}, \quad G_n(t,V) = \sum_{j=1}^{n_j} m_{x,t}(\mathfrak{H}P_j^n)\chi_{P_j^n}(V)$$

where $\chi_B : \mathrm{Grass}(d-l,d) \mapsto \mathbb{R}$ denotes the characteristic function of a measurable set $B \subset \mathrm{Grass}(d-l,d)$. By the Rokhlin disintegration theorem, each $t \mapsto m_{x,t}(\mathfrak{H}P_j^n)$ is a measurable function. It follows that G_n is measurable for every n. Moreover, $(G_n)_n$ converges to G at every point. Thus, G is measurable. Analogously,

$$g_n : K \mapsto \mathbb{R}, \quad g_n(t,V) = \max_{j=1}^{n_j} m_{x,t}(\mathfrak{H}P_j^n)$$

is measurable for every n, and $(g_n)_n$ converges pointwise to g. Thus g is measurable. □

For each fixed $t \in K$, the function $V \mapsto G(t,V)$ is upper semicontinuous: if $(V_n)_n$ converges to V then $\mathfrak{H}V_n$ is contained in a small neighborhood of $\mathfrak{H}V$, for every large n, and then $m_{t,x}(\mathfrak{H}V_n)$ can not be much larger than $m_{t,x}(\mathfrak{H}V)$. Since $\mathrm{Grass}(d-l,d)$ is compact, it follows that the set

$$\Gamma(t) = \{V \in \mathrm{Grass}(d-l,d) : G(t,V) = g(t)\}$$

is compact and non-empty (the supremum in the definition of g is attained) for every $t \in K$.

We borrow the following result from [10, Theorem III.30]:

Theorem 6.3. *Let $(X, \mathfrak{B}, \mu)$ be a complete probability space and Y be a separable complete metric space. Denote by $\mathfrak{B}(Y)$ the Borel $\sigma-$algebra of Y. Let $\kappa(Y)$ be the space of compact subsets of Y, with the Hausdorff topology. The following are equivalent:*

1. *a map $x \to K_x$ from X to $\kappa(Y)$ is measurable;*
2. *its graph $\{(x,y) \in X \times Y : y \in K_x\}$ is in $\mathfrak{B} \otimes \mathfrak{B}(Y)$;*
3. *$\{x \in X : K_x \cap U \neq \emptyset\} \in \mathfrak{B}$ for any open set $U \subset Y$.*

Moreover, any of these conditions implies that there exists a measurable map $\sigma : X \to Y$ such that $\sigma(x) \in K_x$ for every $x \in X$.

Lemma 6.4. *A given $\mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}$ realizes the supremum of*

$$\left\{ m_x(\mathrm{graph}\,\mathfrak{H}\mathcal{V}) : \mathcal{V} \in \mathfrak{H}\,\mathrm{Sec} \right\}$$

if and only if $\mathcal{V}(t) \in \Gamma(t)$ for μ^c-almost every $t \in K$. Moreover, there exists some $\mathcal{V}_x \in \mathfrak{H}\,\mathrm{Sec}$ that does realize this supremum.

Proof. By Lemma 6.2, the set

$$\{(t,V) : V \in \Gamma(t)\} = \{(t,V) : G(t,V) = g(t)\}$$

is a measurable subset of $K \times \mathrm{Grass}(d-l,d)$. Compare the second condition in Theorem 6.3. Thus, from the last claim in the theorem, there exists some measurable map $\mathcal{V}_x : K \to \mathrm{Grass}(d-l,d)$ such that $\mathcal{V}_x(t) \in B_t$ for every $t \in K$. In other words,

$$m_{x,t}(\mathfrak{H}\mathcal{V}_x(t)) = G\big(t, \mathcal{V}_x(t)\big) = g(t) = \sup_Z m_{x,t}(\mathfrak{H}Z)$$

for every $t \in K$. Given any $\mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}$ we have

$$\begin{aligned} m_x(\operatorname{graph}\mathfrak{H}\mathcal{V}) &= \int m_{x,t}(\mathfrak{H}\mathcal{V}(t))\varrho(x,t)\,d\mu^c(t) \\ &\le \int \sup_Z m_{x,t}(\mathfrak{H}Z)\varrho(x,t)\,d\mu^c(t) \\ &= \int m_{x,t}(\mathfrak{H}\mathcal{V}_x(t))\varrho(x,t)\,d\mu^c(t) \\ &= m_x(\operatorname{graph}\mathfrak{H}\mathcal{V}_x). \end{aligned} \tag{24}$$

Thus, $\mathcal{V}_x$ does realize the supremum. Moreover, (24) is an equality if and only if $G(x,\mathcal{V}(t)) = g(t)$ for μ^c-almost every $t \in K$. □

So far, we kept $x \in \Sigma$ fixed. The next proposition shows that the supremum in Lemma 6.4 is actually independent of x. Denote

$$\gamma = \sup\{m_x(\operatorname{graph}\mathfrak{H}\mathcal{V}) : \mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}, x \in \Sigma\}.$$

Proposition 6.5. $\sup\{m_x(\operatorname{graph}\mathfrak{H}\mathcal{V}) : \mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}\} = \gamma$ *for every* $x \in \Sigma$.

Proof. Given any cylinder $[J] \subset \Sigma$, choose a positive constant $c < \mu([J])/\kappa$, where $\kappa > 0$ is the constant in (22). Consider any $\tilde{x} \in \Sigma$ and $\tilde{\mathcal{V}} \in \mathfrak{H}\,\mathrm{Sec}$. For each $k \ge 1$ and $y \in \sigma^{-k}(\tilde{x})$, define $\mathcal{V}^k_y \in \mathfrak{H}\,\mathrm{Sec}$ by

$$\mathcal{F}^k_{\hat{y}}\mathcal{V}^k_y = \tilde{\mathcal{V}}, \text{ that is, } A^k(t,y)\mathcal{V}^k_y(t) = \tilde{\mathcal{V}}(f^k_y(t)) \text{ for each } t \in K.$$

By Corollary 5.11, $m_{\tilde{x}}(\operatorname{graph}\mathfrak{H}\tilde{\mathcal{V}}) = \int m_y(\operatorname{graph}\mathfrak{H}\mathcal{V}^k_y)\,d\mu^u_{k,\tilde{x}}(y)$ and so

$$\begin{aligned} m_{\tilde{x}}(\operatorname{graph}\mathfrak{H}\tilde{\mathcal{V}}) \le{}& \mu^u_{k,\tilde{x}}([J])\sup\{m_x(\operatorname{graph}\mathfrak{H}\mathcal{V}) : \mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}, x \in [J]\} \\ &+ (1 - \mu^u_{k,\tilde{x}}([J]))\gamma. \end{aligned}$$

By (22), there exist arbitrary large values of k such that $\mu^u_{k,\tilde{x}}([J]) \ge c$. Thus

$$\begin{aligned} m_{\tilde{x}}(\operatorname{graph}\mathfrak{H}\tilde{\mathcal{V}}) \le{}& c\sup\{m_x(\operatorname{graph}\mathfrak{H}\mathcal{V}) : \mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}, x \in [J]\}\} \\ &+ (1-c)\gamma. \end{aligned}$$

Varying $\tilde{x} \in \Sigma$ and $\tilde{\mathcal{V}} \in \mathfrak{H}\,\mathrm{Sec}$, we can make the left-hand side arbitrarily close to γ. It follows that

$$\sup\{m_x(\operatorname{graph}\mathfrak{H}\mathcal{V}) : \mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}, x \in [J]\}\} \ge \gamma.$$

The converse inequality is obvious. Thus, we have shown that the supremum over any cylinder $[J]$ coincides with γ.

So, given any $x \in \Sigma$ we may find a sequence $(x_n)_n \to x$ such that the sequence $(m_{x_n}(\operatorname{graph}\mathfrak{H}\mathcal{V}_{x_n}))_n$ converges to γ, where (cf. Lemma 6.4) each $\mathcal{V}_{x_n}$ realizes the supremum at x_n. Moreover, by Proposition 5.5, up to restricting to a subsequence we may assume that $(m_{x_n,t})_n \to m_{x,t}$ for every t in some full μ^c-measure set $X \subset K$. Then

$$\begin{aligned} \gamma &= \lim_n m_{x_n}(\operatorname{graph}\mathfrak{H}\mathcal{V}_{x_n}) \\ &= \lim_n \int m_{x_n,t}(\mathfrak{H}\mathcal{V}_{x_n}(t))\varrho(x_n,t)\,d\mu^c(t) \\ &\le \int \limsup_n m_{x_n,t}(\mathfrak{H}\mathcal{V}_{x_n}(t))\varrho(x_n,t)\,d\mu^c(t). \end{aligned} \tag{25}$$

For each fixed $t \in X$, consider a sub-sequence $(x_{n_k})_k$ along which the lim sup is realized. It is no restriction to suppose that $(\mathcal{V}_{x_{n_k}}(t))_n$ converges to some $V \in \mathrm{Grass}(d-l, d)$ as $k \to \infty$. For any $\epsilon > 0$, let V_ϵ be the closed ϵ-neighborhood of V. The fact that $\mathcal{V}_{x_{n_k}}(t) \subset V_\epsilon$ for every large k implies that

$$\limsup_k m_{x_{n_k},t}(\mathfrak{H}\mathcal{V}_{x_{n_k}}(t)) \le \limsup_k m_{x_{n_k},t}(\mathfrak{H}V_\epsilon) \le m_{x,t}(\mathfrak{H}V_\epsilon)$$

(because V_ϵ is closed). Thus, making $\epsilon \to 0$ on the right-hand side,

$$\limsup_k m_{x_{n_k},t}(\mathfrak{H}\mathcal{V}_{x_{n_k}}(t)) \le m_{x,t}(\mathfrak{H}V) \le m_{x,t}(\mathfrak{H}\mathcal{V}_x(t)).$$

Replacing this in (25), we find that $\gamma \le \int m_{x,t}(\mathfrak{H}\mathcal{V}_x(t))\varrho(x,t)\, d\mu^c(t)$ as claimed. □

Lemma 6.6. *For any $x \in \Sigma$ and $\mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}$, we have $m_x(\mathrm{graph}\,\mathfrak{H}\mathcal{V}) = \gamma$ if and only if $m_y(\mathrm{graph}\,\mathfrak{H}\mathcal{F}_{\hat{y}}^{-1}(\mathcal{V})) = \gamma$ for every $y \in \sigma^{-1}(x)$.*

Proof. Since m is invariant under F, we have

$$m_x = \sum_{\sigma(y)=x} \frac{1}{J\sigma(y)} F_{y*} m_y, \quad \text{with} \quad \sum_{\sigma(y)=x} \frac{1}{J\sigma(y)} = 1,$$

for μ-almost every $x \in \Sigma$. Moreover, since the disintegration $\{m_x : x \in \Sigma\}$ is continuous, the identity extends to every $x \in \Sigma$. Then

$$m_x(\mathrm{graph}\,\mathfrak{H}\mathcal{V}) = \sum_{\sigma(y)=x} \frac{1}{J\sigma(y)} m_y(\mathrm{graph}\,\mathfrak{H}\mathcal{F}_{\hat{y}}^{-1}\mathcal{V}) \tag{26}$$

Since the maximum is γ, it follows from these observations that $m_x(\mathrm{graph}\,\mathfrak{H}\mathcal{V}) = \gamma$ if and only if $m_y(\mathrm{graph}\,\mathfrak{H}\mathcal{F}_{\hat{y}}^{-1}\mathcal{V}) = \gamma$ for every $y \in \sigma^{-1}(x)$. □

As introduced in Section 3.2, let $\hat{\nu} = (\pi_1)_*\hat{\mu}$ and let $\{\hat{\nu}_x : x \in \Sigma\}$ be the disintegration of $\hat{\nu}$ with respect to the partition

$$\{\Sigma^- \times \{x\} : x \in \Sigma\} \sim W^s(x).$$

Observe that every $\hat{\nu}_x$ is equivalent to μ^s.

Lemma 6.7. *For any $x \in \Sigma$ and any $\mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}$ we have that*

$$\hat{m}_{\hat{x}}(\mathrm{graph}\,\mathfrak{H}\mathcal{W}) \le \gamma \text{ for } \hat{\nu}_x \text{ almost every } \hat{x} \in W^s_{loc}(x).$$

Hence, $m_x(\mathrm{graph}\,\mathfrak{H}\mathcal{W}) = \gamma$ if and only if $\hat{m}_{\hat{x}}(\mathrm{graph}\,\mathfrak{H}\mathcal{W}) = \gamma$ for $\hat{\nu}_x$-almost every $\hat{x} \in W^s_{loc}(x)$.

Proof. Suppose there are $y \in \Sigma$,$\mathcal{V} \in \mathfrak{H}\,\mathrm{Sec}$, $\gamma_1 > \gamma$ and a positive $\hat{\nu}_y$-measure subset X of $W^s_{loc}(y)$ such that

$$\hat{m}_{\hat{y}}(\mathrm{graph}\,\mathfrak{H}(\mathcal{V})) \ge \gamma_1 \quad \text{for every } \hat{y} \in X.$$

For each $m \ge 1$, consider the partition of $W^s_{loc}(y) \approx \Sigma^-$ determined by the cylinders $[I]^s = [-m; \iota_{-m}, \dots, \iota_{-1}]^s$, with $\iota_j \in \mathbb{N}$. Since these partitions generate the σ-algebra of $W^s_{loc}(y)$, given any $\varepsilon > 0$ we may find m and I such that

$$\hat{\nu}_y(X \cap [I]^s) \ge (1-\varepsilon)\hat{\nu}_y([I]^s).$$

Note that $[I]^s \times \{y\}$ coincides with $\hat{\sigma}^m(W^s_{loc}(x))$, where $x = \sigma_I^{-m}(y)$. So, using also Lemma 5.2,

$$\hat{\nu}_x\big(\hat{\sigma}^{-m}(X) \cap W^s_{loc}(x)\big) = (\hat{\sigma}^m_*\hat{\nu}_x)(X \cap [I]^s) = J_\mu\sigma^m(x)\, \hat{\nu}_y\big(X \cap [I]^s\big).$$

By the previous inequality and Lemma 5.2, this is bounded below by

$$\begin{aligned}(1-\varepsilon)\, J_\mu \sigma^m(x)\, \hat\nu_y\big([I]^s\big) &= (1-\varepsilon)\, (\hat\sigma_*^m \hat\nu_x)\big([I]^s\big)\\ &= (1-\varepsilon)\, \hat\nu_x\big(W^s_{loc}(x)\big) = 1-\varepsilon.\end{aligned}$$

In this way we have shown that

$$\hat\nu_y\big(\hat\sigma^{-n}(X) \cap W^s_{loc}(x)\big) \ge (1-\varepsilon).$$

Fix $\varepsilon > 0$ small enough so that $(1-\varepsilon)\gamma_1 > \gamma_0$. Using Corollary 5.4, we find that

$$\hat m_{\hat x}(\operatorname{graph} \mathfrak{H}(\mathcal{F}_x^{-n}\mathcal{V})) = \hat m_{\hat y}(\operatorname{graph} \mathfrak{H}(\mathcal{V})) \ge \gamma_1$$

for $\hat\nu_x$-almost every $\hat x \in \hat\sigma^{-n}(X) \cap W^s_{loc}(x)$. It follows that

$$m_x(\operatorname{graph} \mathfrak{H}(\mathcal{F}_x^{-n}\mathcal{V})) = \int \hat m_{\hat x}(\operatorname{graph} \mathfrak{H}(\mathcal{F}_x^{-n}\mathcal{V}))\, d\hat\nu_x(\hat x) \ge (1-\varepsilon)\gamma_1 > \gamma,$$

which contradicts the definition of γ. This contradiction proves the first part of the lemma. The second one is a direct consequence, using

$$m_x(\operatorname{graph} \mathfrak{H}(\mathcal{V})) = \int \hat m_{\hat x}(\operatorname{graph} \mathfrak{H}(\mathcal{V}))\, d\hat\nu_x(\hat x).$$

The proof is complete. □

6.2. **Sections over the periodic point.** Recall (from Section 2.6) that $\hat p$ is a fixed point of $\hat\sigma$ and $\hat z$ is a homoclinic associated to $\hat p$. From now on, we denote $p = P(\hat p)$ and $z = P(\hat z)$. Recall also that $\imath \in \mathbb{N}$ is such that $\hat\sigma^\imath(\hat z) \in W^s_{loc}(\hat p)$.

By the pinching hypothesis in Section 2.6, the Oseledets decomposition of F restricted to K has the form $E^1(t) \oplus \cdots \oplus E^d(t)$, at μ^c-almost every $t \in K$, with $\dim E^i(t) = 1$ for every i. Fix a measurable family $e^1(t), \dots, e^d(t)$ of bases of $\mathbb{C}^d$ with $e^i(t) \in E^i(t)$ for every i. The matrices of the iterates $A^j(p,t)$ relative to these bases are diagonal:

$$A^j(p,t) = \begin{pmatrix} a^{1,j}(t) & 0 & \vdots & 0 \\ 0 & a^{2,j}(t) & \vdots & 0 \\ \cdots & \cdots & \ddots & \cdots \\ 0 & 0 & \vdots & a^{d,j}(t) \end{pmatrix}.$$

We are going to use the associated linear bases of $\Lambda^{(d-l)}\mathbb{C}^d$ and $\Lambda^l\mathbb{C}^d$, defined at μ^c-almost every $t \in K$ by

$$\{e^I(t) = e^{i_1}(t) \wedge \cdots \wedge e^{i_{d-l}}(t), \text{ for } I = \{i_1 < \cdots < i_{d-l}\}\} \tag{27}$$

and

$$\{e^J(t) = e^{j_1}(t) \wedge \cdots \wedge e^{j_l}(t), \text{ for } J = \{j_1 < \cdots < j_l\}\} \tag{28}$$

respectively.

By Lemma 6.4 and Proposition 6.5, we may choose $\mathcal{V}^0 \in \mathfrak{H}\,\mathrm{Sec}$ such that $m_p\big(\operatorname{graph} \mathfrak{H}\, \mathcal{V}^0\big) = \gamma$. Define $\mathcal{V}^j = \mathcal{F}_p^{-j}\mathcal{V}^0$ for $j \ge 1$. By Proposition 6.6, we also have $m_p\big(\operatorname{graph} \mathfrak{H}\, \mathcal{V}^j\big) = \gamma$ for every $j \ge 1$. Let $V^0 : K \to \Lambda^{(d-l)}\mathbb{C}^d$ be a representative of $\mathcal{V}^0 \in \mathfrak{H}\,\mathrm{Sec}$, in the sense that $\mathcal{V}^0(t)$ is the projective class of $V^0(t)$,

for each t. Then denote $V^j = \mathcal{F}_p^{-j} V^0$ for each $j \geq 1$. Expressing V^0 in terms of the linear bases (27) of $\Lambda^{(d-l)}\mathbb{C}^d$,

$$V^0(t) = \sum_{I=i_1,\dots,i_{d-l}} v_I(t) e^I(t),$$

we find that

$$V^j(t) = \sum_{I=i_1,\dots,i_{d-l}} \frac{v_I(f_p^j(t))}{a^{I,j}(t)} e^I(t),$$

with $a^{I,j}(t) = a^{i_1,j}(t) \cdots a^{i_{d-l},j}(t)$. Note that $\lim_j (1/j) \log |a^{i,j}| = \lambda_i$, and so

$$\lim_j \frac{1}{j} \log |a^{I,j}| = \lambda_{i_1} + \cdots + \lambda_{i_{d-l}}. \tag{29}$$

Order the multi-indices

$$I = \{i_1 < \cdots < i_{d-l}\}$$

in such a way that the sums $\lambda_{i_1} + \cdots + \lambda_{i_{d-l}}$ are in increasing order.

Let $\tilde{I}$ be the first multi-index, in this ordering, for which $v_{\tilde{I}}$ is not essentially zero. In what follows we assume that f_p is ergodic for μ^c. Then $\tilde{I}$ is the same for every $t \in K$ in a full μ^c-measure set. The non-ergodic case can be reduced to this one by ergodic decomposition.

Lemma 6.8. *The section $t \mapsto E^{\tilde{I}}(t)$ satisfies $m_p\left(\operatorname{graph} \mathfrak{H} E^{\tilde{I}}\right) = \gamma$.*

Proof. By the Birkhoff ergodic theorem,

$$\lim_n \frac{1}{n} \sum_{j=0}^{n-1} |v_{\tilde{I}}| \left(f_p^j(t)\right) = \int |v_{\tilde{I}}| d\mu_p^c,$$

for μ_p^c-almost every $t \in K$. So, there exist some $\delta > 0$ such that

$$\lim_n \frac{1}{n} \sum_{j=0}^{n-1} |v| \left(f_p^j(t)\right) > \delta > 0$$

for every t in some full μ_p^c-measure set. For any t in that set we may consider a sub-sequence $(j_k)_k$ such that $|v_{\tilde{I}}(f^{j_k q})| > \delta > 0$. Then

$$\lim_k \frac{1}{\|V^{j_k}(t)\|} V^{j_k}(t) = e^{\tilde{I}}(t),$$

and so

$$\lim_k \mathcal{V}^{j_k}(t) = E^{j_1}(t) + E^{j_2}(t) + \cdots + E^{j_l}(t) = E^{\tilde{I}}(t)$$

for μ^c-almost every t. We also have that $m_{p,t}\left(\mathcal{V}^j(t)\right) = \sup_V m_{p,t}(V)$, and then Lemma 6.4 implies that $m_{p,t}(E^{\tilde{I}}(t)) = \sup_V m_{p,t}(V)$ for μ^c-almost every t, as we claimed. □

This means that, from the start, we may take $V^0(t)$ to coincide with one of the invariant sections $E_t^{\tilde{I}}$ given by the Oseledets decomposition, for μ_p^c-almost every $t \in K$. Define $\mathcal{V}' = \mathcal{F}_z^{-\iota} \mathcal{V}^0$. We have that $m_z(\operatorname{graph} \mathfrak{H} \mathcal{V}') = \gamma$ and, by Lemma 6.7, $\hat{m}_{(z^u,z)}(\operatorname{graph} \mathfrak{H} \mathcal{V}') = \gamma$ for μ^s-almost all $(z^u, z) \in W^s_{loc}(\hat{z})$. For each $(x^s, p) \in$

$W^s_{loc}(\hat{p})$, define $\mathcal{V}_{(x^s,p)} = \mathcal{H}^u_{(x^s,z),(x^s,p)}(\mathcal{V}')$, where (x^s, z) is the unique point in $W^u_{loc}((x^s, p)) \cap W^s_{loc}(\hat{z})$. Since $\hat{m}$ is a u-state, and $h^u_{\hat{z},\hat{p}*}\mu^c_{\hat{z}} = \mu^c_{\hat{p}}$, this implies that

$$\hat{m}_{(x^s,p)}(\mathcal{V}_{(x^s,p)}) = \gamma \quad \text{for } \mu^s\text{-almost every } (x^s, p) \in W^s_{loc}(p). \tag{30}$$

Denote $\mathcal{V}^j_{(x^s,p)} = \mathcal{F}^{-j}_{(x^s,p)}\mathcal{V}_{\hat{\sigma}^j((x^s,p))}$ for each $(x^s, p) = \hat{p}$ and $j \geq 1$. In particular, $\mathcal{V}^j_{\hat{p}} = \mathcal{F}^{-j}_{\hat{p}}\mathcal{V}_{\hat{p}}$. We are going to prove that for a large set of js the $\mathcal{V}^j_{\hat{p}}$ have no intersection.

Proposition 6.9. *There exists $N \geq 1$ such that for every $M \in \mathbb{N}$ and $\delta > 0$ there exist $m_1 < m_2 < \cdots < m_M$ and $\tilde{K} \subset K$, with $\mu(\tilde{K}) > 1 - \delta$ and $\mathcal{V}^{m_{k_1}}(t) \cap \cdots \cap \mathcal{V}^{m_{k_N}}(t) = \emptyset$ for any choice of $m_{k_1} < m_{k_2} < \cdots < m_{k_N}$ and $t \in \tilde{K}$.*

Proof. Let $V : K \to \Lambda^{(d-l)}\mathbb{C}^d$ be such that $V(t)$ is an unitary $d - l$ vector that represents $\mathcal{V}_{\hat{p}}(t)$ in $\Lambda^{(d-l)}\mathbb{C}^d$. We can write it as

$$V(t) = \sum_I v_I(t) e^I(t).$$

Then

$$\mathcal{F}^{-j}_{\hat{p}} V(t) = A^j(\hat{p}, t)^{-1} V\left(f^j_p(t)\right) = \sum_I \frac{v_I\left(f^j_p(t)\right)}{a^{I,j}(t)} e^I_t$$

Let $N = \dim \Lambda^l \mathbb{C}^d$. Given any $m_1 < m_2 < \cdots < m_N$ and $t \in K$ such that $\mathcal{V}^{m_1}(t) \cap \cdots \cap \mathcal{V}^{m_N}(t) \neq \emptyset$, there is some non-zero $W(t) \in \Lambda^l \mathbb{C}^d$ such that

$$W(t) \wedge \mathcal{F}^{-m_k}_{\hat{p}} V(t) = 0. \tag{31}$$

Write

$$W(t) = \sum_I w_I(t) e^J_t \quad \text{where } J = \{1, 2, \ldots, l\} \setminus I.$$

Then (31) can be written as

$$\sum_I \frac{v_I\left(f^{m_k}(t)\right)}{a^{J,m_k}(t)} \omega_I(t) \varpi_I = 0,$$

where $\varpi_I = \pm 1$ is the sign of $e^{i_1}_t \wedge \cdots \wedge e^{i_{d-l}}_t \wedge e^{j_1}_t \wedge \cdots \wedge e^{j_{d-l}}_t$. This may be written as

$$B(t)x = 0 \tag{32}$$

where

$$B(t) = \begin{pmatrix} \frac{v_{I_1}(f^{m_1}(t))}{a^{I_1,m_1}(t)} & \cdots & \frac{v_{I_N}(f^{m_1}(t))}{a^{I_N,m_1}(t)} \\ \vdots & \vdots & \vdots \\ \frac{v_{I_1}(f^{m_N}(t))}{a^{I_1,m_N}(t)} & \cdots & \frac{v_{I_N}(f^{m_N}(t))}{a^{I_N,m_N}(t)} \end{pmatrix} \tag{33}$$

$$\text{and } x = \left(\varpi_{I_1}\omega_{I_1}, \ldots, \varpi_{I_N}\omega_{I_N}\right)^T.$$

So, in order to prove that the intersection is necessarily empty, it suffices to show that (32) has no non-zero solutions, in other words, that $\det B(t) \neq 0$. We are going to use the following fact:

Lemma 6.10. *Let $b_i^n : K \to \mathbb{C}$, for $1 \le i \le d$ and $n \in \mathbb{N}$, be measurable functions and suppose there exist $\chi_1 < \chi_2 < \cdots < \chi_d$ such that*

$$\lim_n \frac{1}{n} \log |b_i^n(t)| = \chi_i \text{ for } \mu^c\text{-almost every } t. \tag{34}$$

Then for every $M \in \mathbb{N}$ and $\delta > 0$ there exist $n_1 < n_2 < \cdots < n_M$ and $\tilde{K} \subset K$ with $\mu(\tilde{K}) > 1 - \delta$, such that for any choice of a set $\{k_1, \cdots, k_d\} \subset \{1, \ldots, M\}$ with $k_1 < \cdots < k_d$, the matrix

$$B(t) \in \mathbb{C}^{d \times d}, \quad B_{i,j}(t) = b_i^{n_{k_j}}(t),$$

has non-zero determinant for every $t \in \tilde{K}$

In the proof we are going to use the following simple algebraic fact:

Lemma 6.11. *Let $C = (c_i^j)_{1 \le i,j \le d}$ be a square matrix with $c_i^1 \neq 0$ for every $i = 1, \ldots, d$. Then*

$$\det C = \prod_{i=1}^{d} c_i^1 \cdot \det E$$

where $E = (e_i^j)_{2 \le i,j \le d}$ is defined by

$$e_i^j = \frac{c_i^j}{c_i^1} - \frac{c_1^j}{c_1^1}. \tag{35}$$

Proof. The assumption ensures that we may write

$$\det C = c_1^1 \ldots c_d^1 \begin{bmatrix} 1 & 1 & \ldots & 1 \\ \frac{c_1^2}{c_1^1} & \frac{c_2^2}{c_2^1} & \ldots & \frac{c_d^2}{c_d^1} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{c_1^d}{c_1^1} & \frac{c_2^d}{c_2^1} & \ldots & \frac{c_d^d}{c_d^1} \end{bmatrix}.$$

Subtracting the first column from each one of the others, we end up with

$$\det C = c_1^1 \ldots c_d^1 \begin{bmatrix} e_2^2 & \ldots & e_d^2 \\ \vdots & \ddots & \vdots \\ e_2^d & \ldots & e_d^d \end{bmatrix},$$

as claimed. □

Proof of Lemma 6.10. Let us write $b_i^{1,n} = b_i^n$ for $i = 1, \ldots, d$ and $n \ge 1$. The hypothesis (34) implies that there exist $n_1 \ge 1$ and $K_1 \subset K$ with $\mu(K_1) > 1 - \delta/M$ such that

$$b_i^{1,n}(t) \neq 0 \text{ for } n \ge n_1,\ t \in K_1 \text{ and } i = 1, \ldots, d. \tag{36}$$

Let n_1 be fixed and define (compare (35))

$$b_i^{2,n,n_1}(t) = \frac{b_i^{1,n}(t)}{b_i^{1,n_1}(t)} - \frac{b_1^{1,n}(t)}{b_1^{1,n_1}(t)} \text{ for } i = 2, \ldots, d \text{ and } n > n_1. \tag{37}$$

From (34), and the observation that $\chi_i > \chi_1$, we get that

$$\lim_n \frac{1}{n} \log |b_i^{2,n,n_1}(t)| = \lim_n \frac{1}{n} \log |b_i^{1,n}(t)| = \chi_i. \tag{38}$$

In particular, there exists $n_2 > n_1$ and $K_2 \subset K_1$ with $\mu(K_2) > 1 - 2\delta/M$ such that

$$b_i^{2,n,n_1}(t) \neq 0 \text{ and } b_i^{1,n}(t) \neq 0 \text{ for } n \geq n_2,\ t \in K_2 \text{ and } i = 2, \dots, d \tag{39}$$

(the second condition follows immediately from (36) and the fact that $n_2 > n_1$, but we mention it explicitly, for consistency with what follows).

Next, proceed by induction on $l \leq M$: Suppose that we have defined an increasing sequence of numbers $n_1 < \cdots < n_l$, a decreasing sequence of sets $K_l \supset \cdots \supset K_1$ with $\mu(K_l) > 1 - l\delta/M$, and for every $1 \leq j \leq \max\{l, d\}$ a family of measurable functions $b_i^{j,n_{k_j},\dots,n_{k_1}} : K_l \to \mathbb{C} \setminus \{0\}$ with $1 \leq k_1 < \cdots < k_j \leq l$ satisfying the following relation:

$$b_i^{j,n_{k_j},\dots,n_{k_1}}(t) = \frac{b_i^{j-1,n_{k_j},n_{k_{j-2}},\dots,n_{k_1}}(t)}{b_i^{j-1,n_{k_{j-1}},\dots,n_{k_1}}(t)} - \frac{b_{j-1}^{j-1,n_{k_j},n_{k_{j-2}},\dots,n_{k_1}}(t)}{b_{j-1}^{j-1,n_{k_{j-1}},\dots,n_{k_1}}(t)}, \tag{40}$$

for $i = j, \dots, d$ and $t \in K_l$.

Suppose $l < M$. Fix, $1 < j \leq \max\{l+1, d\}$ and $1 \leq k_1 < \cdots < k_j \leq l$, define the functions $b_i^{j+1,n,n_{k_j},\dots,n_{k_1}} : K_l \to \mathbb{C}$ inductively by

$$b_i^{j+1,n,n_{k_j},\dots,n_{k_1}}(t) = \frac{b_i^{j,n,n_{k_{j-1}},n_{k_{j-2}},\dots,n_{k_1}}(t)}{b_i^{j,n_{k_j},\dots,n_{k_1}}(t)} - \frac{b_j^{j,n,n_{k_{j-1}},n_{k_{j-2}},\dots,n_{k_1}}(t)}{b_j^{j,n_{k_j},\dots,n_{k_1}}(t)} \tag{41}$$

for $i = j+1, \dots, d$, $t \in K_l$ and $n \geq n_l$. Then, arguing as in (38) and using induction on j,

$$\lim_n \frac{1}{n} \log |b_i^{j+1,n,n_{k_j},\dots,n_{k_1}}(t)| = \chi_i. \tag{42}$$

Hence we can find $n_{l+1} > n_l$ and $K_{l+1} \subset K_l$ with

$$\mu(K_{l+1}) > 1 - (l+1)\delta/M$$

such that, for every $1 < j \leq \max\{l+1, d\}$ and $1 \leq k_1 < \cdots < k_j \leq l$,

$$b_i^{j+1,n_{l+1},n_{k_j},\dots,n_{k_1}}(t) \neq 0 \text{ for } i = j+1, \dots, d \text{ and } t \in K_{l+1}.$$

Now fix $\{k_1, \cdots, k_d\} \subset \{1, \dots, M\}$ with $k_1 < \cdots < k_d$, and define for $t \in K_M$ the matrix

$$B(t) \in \mathbb{C}^{d \times d}, \quad B_{i,j}(t) = b_i^{n_{k_j}}(t).$$

Then, in view of the recursive relations (40)–(41), we may apply Lemma 6.11 d-times to $C = B(t)$ to conclude that

$$\det B(t) = \prod_{i=1}^{d} b_i^{1,n_{k_1}}(t) \prod_{i=2}^{d} b_i^{2,n_{k_2},n_{k_1}}(t) \cdots \prod_{i=d}^{d} b_i^{d,n_{k_d},\dots,n_{k_1}}(t).$$

This completes our argument.

□

The twisting condition (Section 2.6), implies that

$$\lim_n \frac{1}{n} \log |v_I(f^n(t))| = 0 \text{ for } \mu^c\text{-almost every } t \in K$$

and $I = \{i_1 < \cdots < i_{d-l}\}$. Then, by (29),

$$\lim_n \frac{1}{n} \log \frac{|v_I(f^n(t))|}{|a^{I,n}(t)|} = -(\lambda_{i_1} + \cdots + \lambda_{i_{d-l}}) \text{ for } \mu^c\text{-almost every } t \in K.$$

The pinching condition ensures that these sums are all distinct. Then we may apply Lemma 6.10 to the functions

$$b_i^n(t) = \frac{v_{I_i}(f^n(t))}{a^{I_i,n}(t)}.$$

We get that there exist $m_1 < \cdots < m_M$ and $\tilde{K} \subset K$ with $\mu(\tilde{K}) > 1 - \delta$ such that for every $\{k_1, \cdots, k_d\} \subset \{1, \ldots, M\}$ with $k_1 < \cdots < k_d$, the matrix $B(t)$ defined in (33) is invertible for every $t \in K_M$. □

Proof of Proposition 6.1. Let $2\delta < \gamma$ and take $C > 0$ large enough that $C(\gamma - 2\delta) > 1$. Consider the sequence of integers $I = \{n_1, n_2, \ldots, n_{CN}\}$ given by Proposition 6.9. Then there exists $\tilde{K} \subset K$ with $\mu^c(K) > 1 - \delta$ such that

$$\mathcal{V}^{n_{k_1}}(t) \cap \cdots \cap \mathcal{V}^{n_{k_N}}(t) = \emptyset \tag{43}$$

for every $t \in \tilde{K}$ and every $\{k_1 < \cdots < k_N\} \subset \{1, \ldots, CN\}$.

First, suppose that $\hat{m}_{\hat{p}}(\operatorname{graph} \mathfrak{H}\mathcal{V}^j_{\hat{p}}) = \gamma$. The property (43) means that the sets $\mathcal{V}^{n_{k_i}}(t)$ are N-wise disjoint for every $t \in \tilde{K}$. Then,

$$\begin{aligned}\hat{m}_{\hat{p}}\Big(\bigcup_{j\in I} \operatorname{graph} \mathfrak{H}\mathcal{V}^j_{\hat{p}}\Big) &\geq \hat{m}_{\hat{p}}\Big(\bigcup_{j\in I} \operatorname{graph} \mathfrak{H}\mathcal{V}^j_{\hat{p}} \mid \tilde{K}\Big) \\ &\geq \frac{1}{N}\sum_I \hat{m}_{\hat{p}}\Big(\operatorname{graph} \mathfrak{H}\mathcal{V}^j_{\hat{p}} \mid \tilde{K}\Big) \geq C(\gamma - \delta) > 1.\end{aligned}$$

This is a contradiction because the measure $\hat{m}_{\hat{p}}$ is a probability.

Now we treat the general case. By (30), $\hat{m}_{(x^s,p)}(\operatorname{graph} \mathfrak{H}\mathcal{V}^{n_j}_{(x^s,p)}) = \gamma$ for every j and μ^s-almost every $(x^s, p) \in W^s_{loc}(p)$. In particular, we may a sequence $\big((x^s_k, p)\big)_k \to \hat{p}$ with that property. Moreover, let $B_k(t)$ be the matrix defined by a system of equations as in (33), with coefficients depending on $\mathcal{V}^{n_i}_{(x^s_k,p)}$ instead of $\mathcal{V}^{n_i}_{\hat{p}}$. Keep in mind that, by definition,

$$\mathcal{V}_{(x^s_k,p)} = \mathcal{H}^u_{(x^s_k,z),(x^s_k,p)}(\mathcal{V}').$$

The sequence $\mathcal{H}^u_{(x^s_k,z),(x^s_k,p)}$ converges uniformly to $\mathcal{H}^u_{\hat{z},\hat{p}}$ when $k \to \infty$. Let $\mathcal{V}_{\hat{p}} = \mathcal{H}^u_{\hat{z},\hat{p}}(\mathcal{V}')$. By Lemma 5.7 (together with the observation that L^1 convergence implies convergence almost everywhere over some subsequence), up to restricting to some subsequence of values of k we have

$$\lim_k \mathcal{V}^{n_i}_{(x^s_k,p)}(t) = \mathcal{V}^{n_i}_{\hat{p}}(t) \text{ for } \mu^c\text{-almost every } t \in K.$$

This proves that B_k converges almost everywhere to B.

Recall that $\det B(t) \neq 0$ for every $t \in \tilde{K}$, by Lemma 6.10. Then, there exist $\tilde{L} \subset \tilde{K}$ with $\mu^c(\tilde{L}) > 1 - 2\delta$ and $k_0 \geq 1$ such that $\det B_k(t) \neq 0$ for every $t \in \tilde{L}$ and $k \geq k_0$. Then, applying the previous argument with (x^s, p) and B_k instead of $\hat{p}$ and B, we get that

$$\begin{aligned}\hat{m}_{(x^s_k,p)}\Big(\bigcup_{j\in I} \operatorname{graph} \mathfrak{H}\mathcal{V}^j_{(x^s_k,p)}\Big) &\geq \frac{1}{N}\sum_I \hat{m}_{(x^s_k,p)}\Big(\operatorname{graph} \mathfrak{H}\mathcal{V}^j_{(x^s_k,p)} \mid \tilde{L}\Big) \\ &\geq C(\gamma - 2\delta) > 1\end{aligned}$$

for every $k \geq k_0$. Thus, again we get a contradiction (because $\hat{m}_{(x^s_k,p)}$ is a probability). □

7. Convergence to Dirac measures

The goal of this section is to prove the following theorem:

Theorem 7.1. *There exists a measurable map $\xi : \hat{M} \to \mathrm{Grass}(l,d)$ such that, given any u-state $\hat{m}$ on $\hat{M} \times \mathrm{Grass}(l,d)$, we have*

$$\hat{m}_{\hat{x},t} = \delta_{\xi(\hat{x},t)} \quad \text{for } \hat{\mu}\text{-almost every } (\hat{x},t) \in \hat{M}.$$

In particular, there exists a unique u-state.

7.1. **Quasi-projective maps.** We begin by recalling the notion of *quasi-projective map*, which was introduced by Furstenberg [12] and extended by Gol'dsheid, Margulis [14]. See also Section 2.3 of [3] for a related discussion.

Let $v \mapsto [v]$ be the canonical projection from $\mathbb{C}^d$ minus the origin to the projective space $\mathbb{PC}^d$. We call $P_\# : \mathbb{PC}^d \to \mathbb{PC}^d$ a *projective map* if there is some $P \in GL(d,\mathbb{C})$ that induces $P_\#$ through $P_\#([v]) = [P(v)]$. The space of projective maps has a natural compactification, the space of *quasi-projective maps*, defined as follows. The quasi-projective map $Q_\#$ induced in $\mathbb{PC}^d$ by a non-zero, possibly non-invertible, linear map $Q : \mathbb{C}^d \to \mathbb{C}^d$ is given by

$$Q_\#([v]) = [Q(v)].$$

Observe that $Q_\#$ is well defined and continuous on the complement of the *kernel* $\ker Q_\# = \{[v] : v \in \ker Q\}$.

More generally, one calls $P_\# : \mathrm{Grass}(l,d) \to \mathrm{Grass}(l,d)$ a *projective map* if there is $P \in GL(d,\mathbb{C})$ that induces $P_\#$ through $P_\#(\xi) = P(\xi)$. Furthermore, the quasi-projective map $Q_\#$ induced in $\mathrm{Grass}(l,d)$ by a non-zero, possibly non-invertible, linear map $Q : \mathbb{C}^d \to \mathbb{C}^d$ is given by

$$Q_\#\xi = Q(\xi).$$

Observe that $Q_\#$ is well defined and continuous on the complement of the *kernel* $\ker Q_\# = \{\xi \in \mathrm{Grass}(l,d) : \xi \cap \ker Q \neq \{0\}\}$.

The space of quasi-projective maps inherits a topology from the space of non-zero linear maps, through the natural projection $Q \mapsto Q_\#$. Clearly, every quasi-projective map $Q_\#$ is induced by some linear map Q such that $\|Q\| = 1$. It follows that the space of quasi-projective maps on any $\mathrm{Grass}(l,d)$ is compact for this topology.

The following two lemmas are borrowed from Section 2.3 of [3]:

Lemma 7.2. *The kernel $\ker Q_\#$ of any quasi-projective map is contained in some hyperplane of $\mathrm{Grass}(l,d)$.*

Lemma 7.3. *If $(P_n)_n$ is a sequence of projective maps converging to some quasi-projective map Q of $\mathrm{Grass}(l,d)$, and $(\nu_n)_n$ is a sequence of probability measures in $\mathrm{Grass}(l,d)$ converging weakly to some probability ν with $\nu(\ker Q) = 0$, then $(P_n)_*\nu_n$ converges weakly to $Q_*\nu$.*

7.2. **Convergence.** Recall that, given $1 \le l \le d-1$ and $1 \le i_1 < \cdots < i_l \le d$, we write

$$E^{i_1,\ldots,i_l}(t) = E^{i_1}(t) \wedge \cdots \wedge E^{i_l}(t) \in \Lambda^l \mathbb{C}^d$$

for every $t \in K$ such that the Oseledets subspaces E^i_t are defined. We give ourselves the right to denote by $E^{i_1,\ldots,i_l}(t)$ also the direct sum

$$E^{i_1}(t) \oplus \cdots \oplus E^{i_l}(t) \in \mathrm{Grass}(l,d).$$

Thus each $E^{i_1,\dots,i_l}$ is an element of $\mathrm{Sec}(K, \mathrm{Grass}(l,d))$.

Let $\hat{p} \in \hat{\Sigma}$ be the fixed point of $\hat{\sigma}$ and $\hat{z} \in \hat{\Sigma}$ be a homoclinic point of $\hat{p}$ with $\hat{z} \in W^u_{loc}(\hat{p})$. Fix $\imath \in \mathbb{N}$ such that $\hat{\sigma}^\imath(\hat{z}) \in W^s_{loc}(\hat{p})$. For each $k \geq 0$, denote $\hat{z}_k = \hat{\sigma}^{-k}(\hat{z})$ and $z_k = P(\hat{z}_k)$. Observe that $\hat{f}_{\hat{z}_k} = f_{z_k}$ and, similarly, $\hat{A}(\hat{p},t) = A(p,t)$. We take advantage of this fact to simplify the notations a bit in the arguments that follow.

Proposition 7.4. *Let $\eta = \mathcal{H}^u_{\hat{p},\hat{z}} E^{1,\dots,l} \in \mathrm{Sec}(K, \mathrm{Grass}(l,d))$. For every sequence $(k_j)_j \to \infty$ there exists a sub-sequence $(k'_i)_i$ such that*

$$\lim_{i\to\infty} A^{k'_i}\big(z_{k'_i}, t_{k'_i}\big)_* m_{z_{k'_i}, t_{k'_i}} = \delta_{\eta(t)}, \text{ where } t_k = (f^k_{z_k})^{-1}(t),$$

for μ^c-almost every $t \in K$.

Proof. We have that $f^k_{z_k} = h^u_{\hat{p},\hat{z}} \circ f^k_p \circ h^u_{\hat{z}_k,\hat{p}}$ and

$$A^k(z_k, t_k) = H^u_{(\hat{p}, h^u_{\hat{z},\hat{p}}(t)),(\hat{z},t)}\, A^k(p, h^u_{\hat{z}_k,\hat{p}}(t_k))\, H^u_{(\hat{z}_k,t_k),(\hat{p}, h^u_{\hat{z}_k,\hat{p}}(t_k))}.$$

So $\big(A^k_{z_k,t_k}\big)_* m_{z_k,t_k}$ is equal to

$$\big(H^u_{(\hat{p}, h^u_{\hat{z},\hat{p}}(t)),(\hat{z},t)}\, A^k(p, h^u_{\hat{z}_k,\hat{p}}(t_k))\big)\big(H^u_{(\hat{z}_k,t_k),(\hat{p}, h^u_{\hat{z}_k,\hat{p}}(t_k))}\big)_* m_{z_k,t_k}.$$

Note that $H^u_{(\hat{z}_k,t_k),(\hat{p}, h^u_{\hat{z}_k,\hat{p}}(t_k))}$ converges uniformly to the identity map id, because $\hat{z}_k$ converges to $\hat{p}$.

Let $K_0 \subset K$ be a full μ^c-measure such that the conclusion of the Oseledets theorem holds at $(\hat{p},t)$ for every $t \in K_0$. We claim that for any $t \in K_0$ and every sub-sequence of

$$A^k(p, h^u_{\hat{z}_k,\hat{p}}(t_k))$$

that converges, the limit is a quasi-projective transformation $Q_\#$ that maps every point outside $\ker Q_\#$ to $E^{1,\dots,l}(h^u_{\hat{z}_k,\hat{p}}(t_k)) \in \mathrm{Grass}(l,d)$. This can be seen as follows.

Given $w \in \Lambda^l \mathbb{C}^d$ and $k \geq 1$, we may write

$$w = \bigoplus_{1\leq i_1<\dots<i_l\leq d} w^{i_1,\dots,i_l}_k E^{i_1,\dots,i_l}\big((f^k_p)^{-1} h_{\hat{z},\hat{p}}(t)\big)$$

with coefficients $w^1_k, \dots, w^N_k \in \mathbb{C}$. It follows from the Oseledets theorem that $k \mapsto w^i_k$ is sub-exponential for every $i = 1, \dots, N$. Recall that $(f^k_p)^{-1}(h(t)) = h_k(t_k)$. Then, the action of $A^k(p, h_k(t_k))$ in the projectivization of the exterior power is given by

$$A^k(p,h_k(t_k))w = \bigoplus_{j=1}^N w^j_k \frac{\Big\| A^k(p, (f^k_p)^{-1}(h(t)) E^{I_j}_{(f^k_p)^{-1}(h(t))} \Big\|}{\big\| A^k(p, (f^k_p)^{-1}(h(t)) \big\|} E^{I_j}_{h(t)}.$$

The quotient of the norms converges to zero for any $j > 1$. Thus, we have that either $A^k(p,h_k(t_k))w \to E^{I_1}_{h(t)}$ or $A^k(p,h_k(t_k))w \to 0$. The latter case means that w is in the kernel of the limit. Thus, any limit quasi-projective transformation does map the complement of the kernel to $E^{I_1}_{h(t)}$, as claimed.

As an immediate consequence we get that for any $t \in K_0$ and every sub-sequence of

$$H^u_{(\hat{p},h(t))),(\hat{z},t)}\, A^k(p, h_k(t_k))$$

that converges, the limit is a quasi-projective transformation that maps every point outside the kernel to $H^u_{(\hat{p},h(t))),(\hat{z},t)} E^{I_1}_{h(t)}$.

By Remark 2.1, the family $\{f^n_{z_k} : n, k \geq 1\}$ is equicontinuous. Using Arzela-Ascoli, it follows that we can find a sub-sequence of $(k_j)_j$ along which the family $(f^k_{z_k})^{-1}$ converges to some $g : K \mapsto K$. Then, by Proposition 10.2, there exists a further subsequence $(k'_i)_i$ and a full μ^c-measure set $K_1 \subset K$ such that

$$m_{z_{k'_i}, t_{k'_i}} \to m_{p,g(t)}$$

for every $t \in K_1$.

By Proposition 6.1 and Lemma 5.8, there exists a full μ^c-measure set $K_2 \subset K$ such that $m_{p,g(t)}$ gives zero weight to every hyperplane of $\mathrm{Grass}(l, d)$ for every $t \in K_2$. Then, by Lemma 7.3 and the previous observations,

$$\lim_{k\to\infty} A^k(z_k, t_k)_* m_{z_k,t_k} = \delta_{\eta(t)}$$

along any sub-sequence such that $A^k(z_k, t_k)$ converges. This yields the claim of the proposition. □

Remark 7.5. The argument remains valid when one replaces the homoclinic point $\hat{z}$ by any other point in $W^u(\hat{p})$.

It follows from Proposition 4.3 that there is a full $\mu^s \times \mu^u$-measure subset of points $\hat{x} \in \hat{\Sigma}$ such that

$$\lim_{n\to\infty} A^n(x_n, t^{\hat{x}}_n)_* m_{x_n, t^{\hat{x}}_n} = \hat{m}_{\hat{x},t} \tag{44}$$

for μ^c-almost every $t \in K$, $x_n = P\big(\hat{\sigma}^{-n}(\hat{x})\big)$ and $t^{\hat{x}}_n = (f^n_{x_n})^{-1}(t)$. Since the shift is ergodic with respect to the projection of $\hat{\mu}$ on $\hat{\Sigma}$, one may also require that

$$\lim_{j\to\infty} \hat{\sigma}^{-n_j}(\hat{x}) = \hat{z}.$$

for some sub-sequence $(n_j)_j \to \infty$.

Fix any $\hat{x} \in \hat{\Sigma}$ such that both conditions hold. Let $k \geq 1$ be fixed, for the time being. Then (44) implies that

$$\begin{aligned} &\lim_{j\to\infty} A^{n_j}(x_{n_j}, t^{\hat{x}}_{n_j})_* m_{x_{n_j}, t^{\hat{x}}_{n_j}} \\ &\qquad = \lim_{j\to\infty} A^{n_j+k}(x_{n_j+k}, t^{\hat{x}}_{n_j+k})_* m_{x_{n_j+k}, t^{\hat{x}}_{n_j+k}} \\ &\qquad = \lim_{j\to\infty} A^{n_j}(x_{n_j}, t^{\hat{x}}_{n_j})_* A^k(x_{n_j+k}, t^{\hat{x}}_{n_j+k})_* m_{x_{n_j+k}, t^{\hat{x}}_{n_j+k}}. \end{aligned} \tag{45}$$

Note also that, by definition,

$$t^{\hat{x}}_{n_j} = f^k_{x_{n_j}+k}(t^{\hat{x}}_{n_j+k}).$$

We use once more the fact that $\{\hat{f}^n_{\hat{x}} : n \in \mathbb{Z} \text{ and } \hat{x} \in \hat{\Sigma}\}$ is equi-continuous (Remark 3.2). Using Ascoli-Arzela, it follows that the exists a sequence $(n_j)_j \to \infty$ such that $(f^{n_j}_{x_{n_j}})^{-1}_j$ converges to some $g : K \to K$. Up to further restricting to a sub-sequence if necessary, Proposition 10.2 ensures that

$$m_{x_{n_j+k}, t^{\hat{x}}_{n_j+k}} \text{ converges to } m_{z_k, g(t)^{\hat{z}}_k} \text{ for } \mu^c\text{-almost every } t.$$

Fix any $t \in K$ such that the previous claims are fulfilled. Let $(n'_i)_i$ be any sub-sequence of $(n_j)_j$ such that $A^{n'_i}\big(x_{n'_i}, t^x_{n'_i}\big)$ converges to some quasi-projective map $Q : \mathrm{Grass}(l, d) \to \mathrm{Grass}(l, d)$. Then (45) may be written as

$$Q_* A^k\big(z_k, g(t)^{\hat{z}}_k\big)_* m_{z_k, g(t)^{\hat{z}}_k}$$

If $\eta(g(t)) \notin \ker Q$ then, making $k \to \infty$, we may use Lemma 7.3 and Proposition 7.4 to conclude that $\hat{m}_{x,t} = \delta_{Q\,\eta(g(t))}$. This gives the conclusion of Theorem 7.1 under this assumption.



FIGURE 1. Proof of Theorem 7.1: avoiding the kernel $\ker Q$

Let us show that we can always reduce the proof to this case. Recall that $\imath \in \mathbb{Z}$ was chosen such that $\hat\sigma^\imath(\hat z) \in W^s_{loc}(\hat p)$. Define $\hat y \in \hat\Sigma$ by

$$\hat\sigma^{n_j+k}(\hat y) \in W^u_{loc}(\hat\sigma^\imath(\hat z)) \cap W^s_{loc}(x_{n_j+k})$$
$$\hat\sigma^\imath(\hat w) \in W^u_{loc}(\hat\sigma^\imath(\hat z)) \cap W^s_{loc}(z_k).$$

Note that $\hat y$ depends on k and j and $\hat w$ that depends on k. We denote $y = P(\hat y)$ and $w = P(\hat w)$. Moreover, $y_n = P(\hat\sigma^{-n}(\hat y))$ and $w_n = P(\hat\sigma^{-n}(\hat w))$ for each $n \geq 0$ Let $m \in \mathbb{N}$ be fixed, for the time being. We have that $x_i = y_i$ with $0 \leq i \leq n_j + k$. So,

$$\hat\sigma^{\imath+m}\big(P(\hat\sigma^{-n_j-k-\imath-m}(\hat y))\big) = y_{n_j+k} = x_{n_j+k}.$$

Also $\hat\sigma^{-n_j-k}(y) \to \hat\sigma^\imath(\hat w)$, and so

$$\hat\sigma^{-n_j-k-\imath-m}(y) \to \hat\sigma^{-m}(\hat w) \text{ when } j \to \infty.$$

Therefore, by Propositions 4.3 and 4.6,

$$\begin{aligned}\hat m_{\hat x,t} &= \lim_{j\to\infty} A^{n_j+k}\big(x_{n_j+k}, t^x_{n_j+k}\big)_* m_{x_{n_j+k}, t^x_{n_j+k}} \\ &= \lim_{j\to\infty} A^{m_j}\big(y_{m_j}, t^{\hat y}_{m_j}\big)_* m_{y_{m_j}, t^{\hat y}_{m_j}}\end{aligned}$$

where $m_j = n_j + k + \imath + m$. The last expression may be rewritten as

$$A^{n_j}\big(x_{n_j}, t^{\hat x}_{n_j}\big)_* A^{k+\imath}\big(y_{n_j+k+\imath}, t^{\hat y}_{y_{n_j+k+\imath}}\big)_* A^m\big(y_{m_j}, t^{\hat y}_{m_j}\big)_* m_{y_{m_j}, t^{\hat y}_{m_j}}.$$

Making $j \to \infty$,

$$\begin{aligned}&\big(f^{n_j+k+\imath}_{y_{n_j+k+\imath}}\big)^{-1} \to \big(f^{k+\imath}_{\hat w}\big)^{-1} \circ g \\ &A^{k+\imath}\big(y_{n_j+k+\imath}, t^{\hat y}_{y_{n_j+k+\imath}}\big) \to A^{k+\imath}\big(w, \big(f^{k+\imath}_w\big)^{-1} g(t)\big) \\ &A^m\big(y_{m_j}, t^{\hat y}_{m_j}\big) \to A^m\big(w_m, \big(f^{k+\imath+m}_{w_m}\big)^{-1} g(t)\big)\end{aligned}$$

and, restricting to a sub-sequence if necessary,

$$m_{y_{m_j},t^{\hat{y}}_{m_j}} \to m_{w_m,\left(f^{k+\imath+m}_{w_m}\right)^{-1}g(t)} \text{ for } \mu^c\text{-almost every } t.$$

Lemma 7.6. *Denote* $\tilde\eta(s) = H^u_{(\hat p,\tilde h(s)),(\hat w,s)} E^{I_1}_{\tilde h(s)}$ *with* $\tilde h(s) = h^u_{\hat w,\hat p}(s)$. *Then there exists a full* μ^c*-measure set* $\tilde K \subset K$ *and a sub-sequence* $(k_j)_j$ *such that for every* $t \in \tilde K$ *there exists a sub-sequence* $(n_i' = n_i'(t))_i$ *of* $(n_j)_j(t)$ *such that*

$$A^{n_i'}\big(x_{n_i'}, t^{\hat x}_{n_i'}\big) \circ A^{k+\imath}\big(y_{n_i'+k_j+\imath}, t^{\hat y}_{y_{n_i'+k_j+\imath}}\big)$$

converges to some quasi-projective transformation $\tilde Q$. *Moreover,* $\tilde\eta\big((f^{k_j+\imath}_w)^{-1}g(t)\big)$ *is not in* $\ker \tilde Q$ *if* j *is sufficiently large, depending on* t.

Proof. As before denote $h = h^u_{\hat z,\hat p}$ and $h_k = h^u_{\hat z_k,\hat p}$. We begin by constructing the sub-sequence $(k_j)_j$. Note that $(f^{k+\imath}_w)^{-1} = \big(f^\imath_w\big)^{-1}\big(f^k_{z_k}\big)^{-1}$ and $\hat w \to \hat z$ when $k \to \infty$. First, take a sub-sequence of values of k such that $\big(f^k_{z_k}\big)^{-1}$ converges uniformly to some ϕ. Since h_k converges to the identity map, $(f^k_p)^{-1}h = h_k(f^k_{z_k})^{-1}$ also converges uniformly to ϕ. Note that ϕ is absolutely continuous with respect to μ^c, by Lemma 5.8. Recall that

$$\tilde\eta\big((f^{k+\imath}_w)^{-1}g(t)\big) = H^u_{(\hat p,\tilde h((f^{k+\imath}_w)^{-1})),(\hat w,(f^{k+\imath}_w)^{-1})} E^{I_1}_{\tilde h((f^{k+\imath}_w)^{-1})}$$

with $\tilde h(s) = h^u_{\hat w,\hat p}(s)$. Up to restricting the sub-sequence of values of k, we may use Lemma 5.7 to get that

$$\begin{aligned} &\tilde\eta\big((f^{k+\imath}_w)^{-1}g(t)\big) \to \eta\big(\big(f^\imath_z\big)^{-1}\phi g(t)\big) \text{ and} \\ &E((f^k_p)^{-1}hg(t)) \to E(\phi g(t))\big) \end{aligned} \tag{46}$$

for every t in some full μ^c-measure set K_1. This defines the sub-sequence $(k_j)_j$ in the statement. In what follows, all the statements on k are meant restricted to this sub-sequence.

The twisting condition implies that

$$\mathcal{H}^u_{\hat z,\hat p}\mathcal{F}^\imath_z E^{I_1}_t \cap \big(E^{j_{l+1}}_t + \cdots + E^{j_d}_t\big) = \{0\} \tag{47}$$

for any $j_{l+1},\dots,j_d \in \{1,\dots,d\}$ and a full μ^c-measure set of values of $t \in K$. In other words, $A^\imath(\eta(h(t))$ does not belong to any of the the hyperplanes of $\mathrm{Grass}(l,d)$ determined by the Oseledets decomposition at the point $(\hat p, t)$. Since ϕ and g are absolutely continuous, there exists a full μ^c-measure set K_2 of values of t such that (47) holds with t replaced by $\phi(g(t))$.

Take $\tilde K = K_1 \cap K_2$. Fix any $t \in \tilde K$ such that in addition $(\hat p, hg(t))$ satisfies the conclusion of the Oseledets theorem. Consider any sub-sequence $(n_i')_i$ of $(n_j)_j$ such that $A^{n_i'}(x_{n_i'}, t^{\hat x}_{n_i'})$ converges to some quasi-projective transformation Q when $i \to \infty$. Then

$$A^{n_i'}\big(x_{n_i'}, t^{\hat x}_{n_i'}\big) \circ A^{k+\imath}\big(y_{n_i'+k+\imath}, t^{\hat y}_{y_{n_i'+k+\imath}}\big)$$

converges to $\tilde Q = Q \circ A^{k+\imath}\big(w, \big(f^{k+\imath}_w\big)^{-1}g(t)\big)$ when $i \to \infty$. Moreover,

$$\begin{aligned} \ker \tilde Q &= A^{k+\imath}\big(w, \big(f^{k+\imath}_w\big)^{-1}g(t)\big)^{-1} \ker Q \\ &= A^\imath\big(w, \big(f^{k+\imath}_w\big)^{-1}g(t)\big)^{-1} A^k\big(z_k, \big(f^k_{z_k}\big)^{-1}g(t)\big)^{-1} \ker Q. \end{aligned}$$

Next, observe that

$$A^k\big(z_k, \big(f^k_{z_k}\big)^{-1} g(t)\big)^{-1} = \Theta_k\, A^{-k}(p, hg(t))\, \Theta \tag{48}$$

where $h = h^u_{\hat{p},\hat{z}}$ and

$$\Theta = H^u_{(\hat{z},g(t)),(\hat{p},hg(t))} \text{ and } \Theta_k = H^u_{(\hat{p},(f^k_p)^{-1}hg(t)),(\hat{z}_k,(f^k_{\hat{z}_k})^{-1}(g(t))}.$$

By Lemma 7.2, the kernel of Q is contained in some hyperplane $\mathfrak{H}v$ of $\mathrm{Grass}(l,d)$. Hence, $\Theta(\ker Q)$ is contained in the hyperplane $\Theta(\mathfrak{H}v)$, of course. Since we take $t \in K$ to be such that the Oseledets theorem holds at $(\hat{p}, t)$, the backward iterates $A^{-k}(p, hg(t))\Theta(\mathfrak{H}v)$ are exponentially asymptotic to some hyperplane section $\mathfrak{H}E$ that is defined by a $(d-l)$-dimensional sum E of Oseledets subspaces. This remains true for $\Theta_k A^{-k}(p, hg(t))\Theta(\mathfrak{H}v)$ because Θ_k converges exponentially fast to to the identity map, since $\hat{z}_k$ converges to $\hat{p}$ exponentially fast. In other words, using (48),

$$\mathrm{dist}_{\mathrm{Grass}(l,d)}\left(A^k\big(z_k, \big(f^k_{z_k}\big)^{-1} g(t)\big)^{-1}\mathfrak{H}v, \mathfrak{H}E((f^k_p)^{-1}hg(t))\right) \to 0$$

exponentially fast as $k \to \infty$. Then, by (46), we have that $A^k\big(z_k, \big(f^k_{z_k}\big)^{-1} g(t)\big)^{-1}\mathfrak{H}v$ converges to $E(\phi g(t))$. So,

$$\ker \tilde{Q} \subset A^\imath\big(z, (f^\imath_z)^{-1}\phi g(t)\big)^{-1}\mathfrak{H}E(\phi g(t)).$$

Keep in mind that $\hat{z} \in W^u_{loc}(\hat{p})$ and $\imath \in \mathbb{N}$ is such that $\hat{\sigma}^\imath(\hat{z}) \in W^s_{loc}(\hat{p})$. Recall also (from Section 4.1) that in the present setting all the local stable holonomies h^s and H^s are trivial. In particular, (8) means that

$$V^i(t') = H^u_{(\hat{z},t_1),(\hat{p},t')}\hat{A}^{-\imath}(\hat{\sigma}^\imath(\hat{z}), s)E^i(s) = H^u_{(\hat{z},t_1),(\hat{p},t')}\hat{A}^\imath(\hat{z}, t_1)^{-1}E^i(s)$$

with $t' = \phi g(t)$, $t_1 = h^u_{\hat{p},\hat{z}}(t')$ and $s = \hat{f}^\imath_{\hat{z}}(t_1)$. Then, by the twisting condition, $\oplus_{j\in J}V^j$ cannot intersect any sum of the form $\oplus_{i\in I}E^i$ with $\#I + \#J = d$. In particular, the distance between

$$H^u_{(\hat{z},t_1),(\hat{p},t')}\hat{A}^{-\imath}(\hat{\sigma}^\imath(\hat{z}), s)E(s) \quad \text{and} \quad E^{I_1}(s)$$

is positive. Equivalently, the distance between

$$\hat{A}^{-\imath}(\hat{\sigma}^\imath(\hat{z}), s)E(s) \quad \text{and} \quad \eta(t_1) = H^u_{(\hat{p},t'),(\hat{z},t_1)}E^{I_1}(s)$$

is positive. Then $\eta\big(\big(f^\imath_z\big)^{-1}\phi g(t)\big)$ does not intersect

$$\hat{A}^{-\imath}(\hat{\sigma}^\imath(\hat{z}), \big(\phi g(t)\big))E\big(\phi g(t)\big) = A^\imath\big(z, (f^\imath_z)^{-1}\phi g(t)\big)^{-1}E(\phi g(t)),$$

which implies that $\eta\big(\big(f^\imath_z\big)^{-1}\phi g(t)\big) \notin \ker \tilde{Q}$. This gives the conclusion of the lemma. □

Having established Lemma 7.6, we can now use the same argument as previously, to conclude that $\hat{m}_{\hat{x},t} = \delta_{\tilde{Q}\eta}$ at μ^c-almost every point also in this case. To do this, observe that for every m and k fixed there exist a sub-sequence $(m'_i)_i$ of $(m_j)_j$ such that

$$m_{y_{m'_i}, t^{\hat{y}}_{m'_i}} \to m_{w_m, \left(f^{k+\imath+m}_{w_m}\right)^{-1} g(t)} \text{ for } \mu^c\text{-almost every } t. \tag{49}$$

Using a diagonal argument, we may choose $(m'_i)_i$ to be independent of k and m. Fix, once and for all, a full μ^c-measure subset K' such that (49) and the conclusions of Lemma 7.6 and Proposition 7.4 (more precisely, Remark 7.5) hold for every $t \in K'$.

For each fixed $t \in K'$, fixing k sufficiently large and making m_i' go to infinity (along the sub-sequence given by Lemma 7.6), we find that

$$\hat{m}_{\hat{x},t} = \tilde{Q}_*\big(A^m(w_m, (f_{w_m}^{k+\iota})^{-1} g(t)\big)_* m_{w_m,(f_{w_m}^{k+\iota})^{-1}g(t)}$$

Then, making $m \to \infty$ and using Lemma 7.3 and Proposition 7.4,

$$\hat{m}_{\hat{x},t} = \delta_{\xi(\hat{x},t)},$$

where $\xi(\hat{x},t) = \tilde{Q}\tilde{\eta}((f_w^{k+\iota})^{-1} g(t)$.

Thus we proved that $\hat{m}_{\hat{x},t}$ is a Dirac measure for $\hat{\nu}$-almost every $\hat{x} \in \hat{\Sigma}$ and $\hat{\mu}_{\hat{x}}^c$-almost every $t \in K$. Note also that the set $\tilde{M} \subset \hat{M}$ of points $(\hat{x},t) \in \hat{M}$ such that $\hat{m}_{(\hat{x},t)}$ is a Dirac measure is measurable, since the map $(\hat{x},t) \mapsto \hat{m}_{(\hat{x},t)}$ is measurable and the set of Dirac measures is closed in the weak* topology is closed, then $\tilde{M}$ is measurable. Thus we have shown that $\tilde{M}$ has total $\hat{\mu}$-measure, which completes the proof of Theorem 7.1.

8. Orthogonal complement

8.1. **Eccentricity.** Let $L : \mathbb{C}^d \to \mathbb{C}^d$ be a linear isomorphism and $1 \le l \le d$. The l-dimensional *eccentricity* of L is defined by

$$E(l, L) = \sup\Big\{\frac{m(L \mid \xi)}{\|L \mid \xi^\perp\|} : \xi \in \mathrm{Grass}(l,d)\Big\}, \quad m(L \mid \xi) = \big\|(L \mid \xi)^{-1}\big\|^{-1}.$$

We call *most expanded* l-subspace any $\xi \in \mathrm{Grass}(l,d)$ that realizes the supremum. These always exist, since the Grassmannian is compact and the expression depends continuously on ξ.

These notions may be expressed in terms of the polar decomposition of $L = K'DK$ with respect to any orthonormal basis: denoting by $a_1, \dots, a_d$ the eigenvalues of the diagonal operator D, in non-increasing order, then $E(l,L) = a_l/a_{l+1}$. The supremum is realized by any subspace ξ whose image under K is a sum of l-eigenspaces of D such that the product of the eigenvalues is $a_1 \cdots a_l$. It follows that $E(l,L) \ge 1$, and the most expanded l-subspace is unique if and only if the eccentricity is strictly larger than 1.

Proposition 8.1. *For every $0 < c < 1$, there exists a set $\hat{M}_c \subset \hat{M}$ with $\hat{\mu}(\hat{M}_c) > c$ such that $E\big(l, A^n(\hat{f}^{-n}(\hat{x},t))\big) \to \infty$, and the image of the most expanded subspace by $A^n(\hat{f}^{-n}(\hat{x},t))$ converges to $\xi(\hat{x},t)$, restricted to the iterates such that $\hat{f}^{-n}(\hat{x},t) \in \hat{M}_c$*

For the proof, let us recall the following proposition, whose proof can be found in [3]:

Proposition 8.2. *Let $\mathcal{N}$ be a weak* compact family of probabilities on $\mathrm{Grass}(l,d)$ such that all $\nu \in \mathcal{N}$ give zero weight to every hyperplane. Let $L_n : \mathbb{C}^d \to \mathbb{C}^d$ be linear isomorphisms such that $(L_n)\nu_n$ converges to a Dirac measure δ_ξ as $n \to \infty$, for some sequence ν_n in N. Then the eccentricity $E(l, L_n)$ goes to infinity and the image $L_n(\zeta_n)$ of the most expanding l-subspace of L_n converges to ξ.*

Proof of Proposition 8.1. Given $0 < c < 1$ take $M_c \subset M$ to be a compact set, with $\mu(M_c) > c$ and such that the restriction of the map $(x,t) \mapsto m_{(x,t)}$ to M_c is continuous. This implies that

$$\mathcal{N} = \{m_{(x,t)}; (x,t) \in M_c\}$$

is a weak* compact subset of the space of probability measures of Grass(l,d), and every measure in $\mathcal{N}$ gives zero weight to every hyperplane. Moreover,

$$A^n(\hat f^{-n}(\hat x,t))_* m_{P\times \mathrm{id}(\hat f^{-n}(\hat x,t))} = \delta_{\xi(\hat x,t)}.$$

Take $\hat M_c = (P\times \mathrm{id})^{-1}(M_c)$. Then the claim follows from Proposition 8.2, with $L_n = A^n(\hat f^{-n}(\hat x,t))$. □

8.2. **Adjoint cocycle.** Fix a continuous Hermitian form $\langle\cdot,\cdot\rangle_{(\hat x,t)}$ in $\hat M\times\mathbb{C}^d$. Let $\hat F^*:\hat M\times\mathbb{C}^d\to\hat M\times\mathbb{C}^d$ be the adjoint cocycle, defined over $\hat f^{-1}:\hat M\to\hat M$ by

$$\hat F^*((\hat x,t),v) = (\hat f^{-1}(\hat x,t), \hat A_*(\hat x,t)v)$$

where $\hat A_*(\hat x,t)$ is the adjoint $\hat A(\hat f^{-1}(\hat x,t))^*$ of the matrix $\hat A(\hat f^{-1}(\hat x,t))$ with respect to the Hermitian form. In other words, $\hat A_*(\hat x,t)$ is characterized by

$$\Big\langle u, \hat A_*(\hat x,t)v\Big\rangle_{\hat f^{-1}(\hat x,t)} = \Big\langle \hat A(\hat f^{-1}(\hat x,t)u), v\Big\rangle_{(\hat x,t)} \quad \text{for any } u,v\in\mathbb{C}^d$$

and $(\hat x,t)\in\hat M$. We have that

$$W^{ss}_{\hat f^{-1}}(\hat x,t) = W^{uu}_{\hat f}(\hat x,t) \quad\text{and}\quad W^{uu}_{\hat f^{-1}}(\hat x,t) = W^{ss}_{\hat f}(\hat x,t).$$

It is also easy to see that

$$H^{u,\hat A_*}_{(\hat x,t),(\hat y,s)} = (H^{s,\hat A}_{(\hat y,s),(\hat x,t)})^* \text{ and } H^{s,\hat A_*}_{(\hat x,t),(\hat z,r)} = (H^{u,\hat A}_{((\hat z,r)),(\hat x,t)})^*,$$

respectively, for any $(\hat x,t),(\hat y,s)$ in the same $\hat f^{-1}$-unstable set and any $(\hat x,t),(\hat z,r)$ in the same $\hat f^{-1}$-stable set.

The following result is probably well-known but we include a proof since we could not find one in the literature:

Proposition 8.3. *The cocycles $\hat F$ and $\hat F^*$ have the same Lyapunov exponents. Moreover, if E^j, $j=1,\dots,k$ are the Oseledets spaces of $\hat F$ then the Oseledets spaces of $\hat F^*$ are, respectively,*

$$E^j_* = \big[E^1\oplus\cdots\oplus E^{j-1}\oplus E^{j+1}\oplus\cdots\oplus E^k\big]^\perp, j=1,\dots,k.$$

Proof. (We restrict ourselves to the case when $\dim E^j = 1$ for every j, which is the only case we will actually use; the extension to the general case is rather straightforward.)

First of all, if $e^j_*(x)\in E^j_*(x)$ and $v^j\in E^j_*(f^{-1}(x))^\perp$ then

$$\Big\langle v^j, \hat A_*(x)e^j_*(x)\Big\rangle = \Big\langle \hat A(f^{-1}(x))v^j, e^j_*(x)\Big\rangle = 0$$

and so $\hat A_*(x)E^j_*(x) = E^j_*(f^{-1}(x))$. This shows that the decomposition $E^1_*\oplus\cdots\oplus E^k_*$ is invariant under $\hat A_*$.

Let $e^j_*(x)\in E^j_*(x)$ and $e^j(x)\in E^j(x)$ be unit vectors. Then

$$\Big\langle e^k(f^{-n}(x)), \hat A^n_*(x)e^j_*(x)\Big\rangle = 0$$

for $k\neq j$, and

$$\Big\langle e^j(f^{-n}(x)), \hat A^n_*(x)e^j_*(x)\Big\rangle = \Big\|\hat A^n_*(x)e^j_*(x)\Big\| \cos(\alpha^j(f^{-n}(x)), \tag{50}$$

where

$$\begin{aligned}\alpha^j(x) &= \measuredangle\left(e^j_*(x), e^j(x)\right)\\ &= \frac{\pi}{2} - \measuredangle\left(e^j_*(x), E^1_x \oplus \cdots \oplus \hat{E^j}_x \oplus \cdots \oplus E^k_x\right).\end{aligned}$$

So, by the Oseledets theorem,

$$\lim_{n\to\infty} \frac{1}{n} \log|\cos(\alpha^j(f^{-n}(x))| = 0.$$

On the other hand,

$$\begin{aligned}&\left\langle A^n(f^n(x))e^j(f^{-n}(x)), e^j_*(x)\right\rangle\\ &\qquad = \left\|A^n(f^n(x))e^j(f^{-n}(x))\right\| \cos(\alpha^j(x)).\end{aligned} \tag{51}$$

Moreover, by the invariance of e^j,

$$\frac{A^n(f^n(x))e^j(f^{-n}(x))}{\|A^n(f^n(x))e^j(f^{-n}(x))\|} = e^j(x).$$

Thus

$$\begin{aligned}-\lambda_j &= \lim_{n\to\infty} \frac{1}{n} \log\left\|A^{-n}(x)e^j(x)\right\|\\ &= \lim_{n\to\infty} \frac{1}{n} \log\left\|\frac{e^j(f^{-n}(x))}{\|A^n(f^n(x))e^j(f^{-n}(x))\|}\right\|\\ &= -\lim_{n\to\infty} \frac{1}{n} \log\left\|A^n(f^n(x))e^j(f^{-n}(x))\right\|.\end{aligned} \tag{52}$$

Putting (50), (51) and (52) together we get that

$$\lim_{n\to\infty} \frac{1}{n}\left\|\hat{A}^n_*(x)e^j_*(x)\right\| = \lambda_j,$$

which proves the statement. □

Proposition 8.4. *$\hat{A}$ is simple, if and only if, $\hat{A}_*$ is simple.*

Proof. Applying Proposition 8.3 to the restriction of $\hat{F}$ to the periodic leaf $\{\hat{p}\}\times K$ we get that the cocycle $\hat{A}_*$ is pinching, if and only if, $\hat{A}$ is pinching. Moreover, the Oseledets decomposition $E^1_*\oplus\cdots\oplus E^k_*$ of $\hat{A}_*$ is given by the orthogonal complements of the Oseledets subspaces of A:

$$E^j_* = \left[E^1 \oplus \cdots \oplus E^{j-1} \oplus E^{j+1} \oplus \cdots \oplus E^k\right]^\perp.$$

We are going to use this for proving the twisting property, as follows.

Let $\phi_{\hat{p},\hat{z}} = \mathcal{H}^{s,\hat{A}}_{\hat{z},\hat{p}} \circ \mathcal{H}^{u,\hat{A}}_{\hat{p},\hat{z}}$ and $\phi^*_{\hat{p},\hat{z}} = \mathcal{H}^{s,\hat{A}_*}_{\hat{z},\hat{p}} \circ \mathcal{H}^{u,\hat{A}_*}_{\hat{p},\hat{z}}$. Denote

$$\begin{aligned}&h : K \to K, \quad h(t) = h^s_{\hat{z},\hat{p}} \circ h^u_{\hat{p},\hat{z}} \quad \text{and}\\ &H_t = H^s_{(\hat{z},h^s_{\hat{p},\hat{z}}(t)),(\hat{p},t)} \circ H^u_{(\hat{p},h^{-1}(t)),(\hat{z},h^s_{\hat{p},\hat{z}}(t))},\end{aligned}$$

Then, for any $V \in \mathrm{Sec}(K, \mathrm{Grass}(l,d))$,

$$\phi_{\hat{p},\hat{z}}V(t) = H_t\left(V(h^{-1}(t)\right) \quad \text{and} \quad \phi^*_{\hat{p},\hat{z}}V(t) = {H_{h(t)}}^*\left(V(h(t)\right).$$

First, we treat the case $l = 1$. Write $g = f_{\hat{p}}$ and let $e^j(t) \in E^j(t)$ and $e^j_*(t) \in E^j_*(t)$ be unit vectors. The twisting condition means that if

$$\phi_{\hat{p},\hat{z}} e^k(t) = \sum_{j=1}^{d} a_{k,j}(t) e^j(t),$$

then

$$\lim_{n\to\infty} \frac{1}{n} \log|a_{k,j}(g^n(t))| = 0.$$

We need to deduce the corresponding fact for the adjoint. For this, write

$$\phi^*_{\hat{p},\hat{z}} e^k_*(t) = \sum_{j=1}^{d} b_{k,j}(t) e^j_*(t).$$

Hence

$$\begin{aligned} \beta_{k,j}(t)\langle e^j_*(t), e^j(t)\rangle &= \langle \phi^*_{\hat{p},\hat{z}} e^k_*(t), e^j(t)\rangle \\ &= \langle e^k_*(h(t)), \phi_{\hat{p},\hat{z}} e^j(h(t))\rangle \\ &= \overline{a_{j,k}(h(t))}\langle e^k_*(h(t)), e^k(h(t))\rangle, \end{aligned}$$

by definition $\langle e^i_*(t), e^i(t)\rangle = \cos(\alpha^i(x))$ for every $1 \le i \le d$, where

$$\begin{aligned} \alpha^i(t) &= \measuredangle\left(e^j_*(t), e^j(t)\right) \\ &= \frac{\pi}{2} - \measuredangle\left(e^j_*(t), E^1_t \oplus \cdots \oplus \hat{E^j}_t \oplus \cdots \oplus E^k_t\right), \end{aligned}$$

so, by the Oseledets theorem,

$$\lim_{n\to\infty} \frac{1}{n} \log|\langle e^i_*(g^n(t)), e^i(g^n(t))\rangle| = 0 \ \hat{\mu}^c_{\hat{p}}\text{-almost everywhere.}$$

Then

$$\lim_{n\to\infty} \frac{1}{n} \log|\beta_{k,j}(g^n(t))| = \lim_{n\to\infty} \frac{1}{n} \log|a_{j,k}(g^n(h(t)))|$$

also as $h : K \to K$ preserves $\hat{\mu}^c_{\hat{p}}$ this is true for $\hat{\mu}^c_{\hat{p}}$-almost everywhere.

For $l > 1$ the proof is exactly the same, using the inner product induced on $\Lambda^l\mathbb{C}^d$ by $\langle\cdot,\cdot\rangle$, that is,

$$\langle v_1 \wedge \cdots \wedge v_l, w_1 \wedge \cdots \wedge w_l\rangle_{\Lambda^l\mathbb{C}^d} = \det\left(\langle v_i, w_j\rangle\right).$$

Thus, we have shown that A is twisting if and only if A_* is twisting. □

Applying Proposition 8.1 to the adjoint cocycle we get:

Corollary 8.5. *There exists a section $\xi^* : \hat{M} \to Grass\,(l,d)$ which is invariant under the cocycle $F_{\hat{A}_*}$ and the unstable holonomies of $\hat{A}_*$.*

Moreover, given any $c > 0$ there exists $\hat{M}_c \subset \hat{M}$ with $\hat{\mu}(\hat{M}_c) > c$ such that, restricted to the sub-sequence of iterates k such that $\hat{f}^k(p)$ in $\hat{M}_c$, the eccentricity $E\big(l, \hat{A}^k_*(\hat{f}^k(p))\big) = E\big(l, A^k(p)\big)$ goes to infinity and the image $\hat{A}^k_*\big(\hat{f}^k(p)\big)\zeta^a_k(\hat{f}^k(p))$ of the most expanded l-subspace tends to $\xi^*(p)$ as $k \to \infty$.

The next lemma relates the invariant sections of the two cocycles, F and $F_{\hat{A}_*}$:

Lemma 8.6. *For $\hat{\mu}$-almost every $\hat{x}, t \subset \hat{M}$, the subspace $\xi(\hat{x}, t)$ is transverse to the orthogonal complement of $\xi^*(\hat{x}, t)$.*

Proof. Recall that the stable holonomies of $\hat{A}$ are trivial. Thus, the unstable holonomies of $\hat{A}_*$ are trivial. So, the fact that ξ^* is invariant under unstable holonomies means that it is constant on local stable sets of $\hat{f}$. Then the same is true about his orthogonal complement $\eta(\hat{x},t) = \xi^*(\hat{x},t)^\perp$, which means that it only depends on $\eta(\hat{x},t) = \eta(x,t)$, where $x = P(\hat{x})$. So the graph of $\eta(x,\cdot)$ over K has zero m_x-measure:

$$\begin{aligned} m_x(\operatorname{graph}\mathfrak{H}\eta_x) &= \int\int \delta_{\xi_{\hat{x},t}}(\eta(x,t))d\mu^c(t)d\mu^s_x(\hat{x}) \\ &= \mu^c \times \mu^s\left(\{\hat{x},t : \xi(\hat{x},t) \in \eta(x,t)\}\right) \\ &= 0 \end{aligned}$$

for ν-almost every $x \in \Sigma$. Hence $\hat{\mu}\left(\{\hat{x},t : \xi(\hat{x},t) \in \eta(x,t)\}\right) = 0$, which proves the lemma. □

9. Proof of Theorem B

Let $(\hat{x},t) \in \hat{M}$, and denote by $\eta(\hat{x},t) \in Grass\,(d-l,d)$ the orthogonal complement of $\xi^*(\hat{x},t)$. Recall that ξ^* is invariant under $\hat{A}_*$, in the sense that

$$\hat{A}_*(\hat{x},t)\xi^*(\hat{x},t) = \xi^*(\hat{f}^{-1}(\hat{x},t)).$$

Hence, η is invariant under A.

According to Lemma 8.6, we have that $\mathbb{C}^d = \xi(\hat{x},t) \oplus \eta(\hat{x},t)$ at $\hat{\mu}$-almost every point. We want to prove that the Lyapunov exponents of A along ξ are strictly bigger than those along η. For that, let

$$\xi(\hat{x},t) = \xi^1(\hat{x},t) \oplus \cdots \oplus \xi^u(\hat{x},t) \quad \text{and} \quad \eta(\hat{x},t) = \eta^s(\hat{x},t) \oplus \cdots \oplus \eta^1(\hat{x},t)$$

be the Oseledets decomposition of A restricted to the two invariant sub-bundles, where ξ^u corresponds to the smallest Lyapunov exponent among ξ^i and η^s the largest among all η^j.

Denote $d_u = \dim \xi^u$ and $d_s = \dim \eta^s$, and then let λ_u and λ_s be the Lyapunov exponents associated to these two sub-bundles, respectively. Define

$$\Delta^n(\hat{x},t) = \frac{\det\left(A^n(\hat{x},t), \xi^u(\hat{x},t)\right)^{\frac{1}{d_u}}}{\det\left(A^n(\hat{x},t), W(\hat{x},t)\right)^{\frac{1}{d_u+d_s}}},$$

where $W(\hat{x},t) = \xi^u(\hat{x},t) \oplus \eta^s(\hat{x},t)$. By the Oseledets theorem

$$\lim_{n\to\infty} \frac{1}{n} \log \Delta^n(\hat{x},t) = \frac{d_s}{d_u+d_s}\left(\lambda_u - \lambda_s\right).$$

The proof of the following proposition is identical to the proof of Proposition 7.3 in [3]:

Proposition 9.1. *For every $0 < c < 1$ there exist a set $\hat{M}_c \subset \hat{M}$ with $\hat{\mu}(\hat{M}_c) > c$ such that for $\hat{\mu}$-almost every $(\hat{x},t) \in \hat{M}$*

$$\lim_{n\to\infty} \Delta^n(\hat{x},t) = \infty$$

restricted to the sub-sequence of values n for which $\hat{f}^n(\hat{x},t) \in \hat{M}_c$.

So now fix some $0 < c < 1$ and $\hat{M}_c$ given by Proposition 9.1. Let $g : \hat{M}_c \to \hat{M}_c$ be the first return map:

$$g\left(\hat{x},t\right) = \hat{f}^{r(\hat{x},t)}\left(\hat{x},t\right).$$

Then we can define the induced cocycle $G : \hat{M}_c \times \mathbb{C}^d \to \hat{M}_c \times \mathbb{C}^d$

$$G\left((\hat{x},t),v\right) = \left(g\left(\hat{x},t\right), D(\hat{x},t)v\right),$$

where $D(\hat{x},t) = \hat{A}^{r(\hat{x},t)}\left(\hat{x},t\right)$. It is well-known (see [20, Proposition 4.18]) that the Lyapunov exponents of G with respect to $\frac{1}{\hat{\mu}(\hat{M}_c)}\hat{\mu}$ are the products of the exponents of $\hat{F}_A$ by the average return time $1/\hat{\mu}(\hat{M}_c)$. Thus, to show that $\lambda_u > \lambda_s$ it suffices to prove the corresponding fact for G.

Define

$$\tilde{\Delta}^k((\hat{x},t)) = \frac{\det\left(D^k(\hat{x},t), \xi^u(\hat{x},t)\right)^{\frac{1}{d_u}}}{\det\left(D^k(\hat{x},t), W(\hat{x},t)\right)^{\frac{1}{d_u+d_s}}}.$$

Then $\tilde{\Delta}^k(\hat{x},t)$ is a sub-sequence of $\Delta^n(\hat{x},t)$ such that $\hat{f}^n(\hat{x},t) \in \hat{M}_c$. So, using Proposition 9.1 we conclude that

$$\lim_{n\to\infty}\sum_{j=0}^{k-1}\log\tilde{\Delta}\left(g^j(\hat{x},t)\right) = \lim_{n\to\infty}\log\tilde{\Delta}^k\left(p\right) = \infty$$

for $\hat{\mu}$-almost every $(\hat{x},t) \in \hat{M}_c$.

We need the following well-known fact:

Lemma 9.2. *Let $T : X \to X$ be a measurable transformation preserving a probability measure ν in X, and $\varphi : X \to \mathbb{R}$ be a ν integrable function such that $\lim_{n\to\infty}\sum_{j=0}^{n-1}\left(\varphi\circ T^j\right) = +\infty$ at ν almost every point. Then $\int\varphi d\nu > 0$.*

Applying the lemma to $T = g$ and $\varphi = \log\tilde{\Delta}$ we find that

$$\lim_{k\to\infty}\frac{1}{k}\log\tilde{\Delta}^k(\hat{x},t) = \lim_{k\to\infty}\frac{1}{k}\sum_{j=0}^{k-1}\log\tilde{\Delta}\left(g^j(\hat{x},t)\right) = \int\log\tilde{\Delta}\frac{d\hat{\mu}}{\hat{\mu}(\hat{M}_c)} > 0$$

at $\hat{\mu}$-almost every point. On the other hand the relation between Lyapunov exponents gives that

$$\lim_{k\to\infty}\frac{1}{k}\log\tilde{\Delta}^k(\hat{x},t) = \frac{d_s}{d_u+d_s}\left(\lambda_u-\lambda_s\right)\frac{1}{\hat{\mu}(\hat{M}_c)}.$$

this means that $\lambda_u > \lambda_s$, so there is a gap between the first l Lyapunov exponents and the remaining $d-l$ ones. Since this applies for every $1 \le l \le d$, we conclude that the Lyapunov spectrum is simple. This completes the proof of Theorem B.

10. Proof of Theorem C

We begin by proving

Lemma 10.1. *The set of uniformly simple cocycles is non-empty.*

Proof. Choose $A \in GL(d,\mathbb{C})$ such that its eigenvalues $\tau_1,\ldots,\tau_d$ satisfy

$$\tau_1^{m+1}\cdots\tau_d^{m_d} = 1 \text{ with } m_1,\ldots,m_d \in \mathbb{Z} \text{ only if } m_1 = \cdots = m_d = 0.$$

In particular, the eigenvalues $\tau_1,\ldots,\tau_d$ are all distinct. Then let $\hat{A} : \hat{M} \to GL(d,\mathbb{C})$ be the constant cocycle $\hat{A}(\hat{x},t) \equiv A$. It is clear that $\hat{A}$ admits (trivial) stable and unstable holonomies: $H^*_{\cdot,\cdot} = \mathrm{id}$ for every $* \in \{s,u\}$.

Let us call $h^u(t) = h^u_{\hat{p},\hat{z}}(t)$ and $h = h^s_{\hat{z},\hat{p}}\circ h^u_{\hat{p},\hat{z}}$, and denote

$$H^{\hat{A}}_t = H^{s,\hat{A}}_{(\hat{z},h^u(t)),(\hat{p},h(t))}H^{u,\hat{A}}_{(\hat{p},t),(\hat{z},h^u(t))}.$$

By the definition of holonomies,

$$H^{s,\hat{A}}_{(\hat{z},h^u(t)),(\hat{p},h(t))} = \hat{A}(\hat{p},h(t))^{-1} H^{s,\hat{A}}_{\hat{\sigma}(\hat{z},h^u(t)),\hat{\sigma}(\hat{p},h(t))} \hat{A}(\hat{x},h^u(t)).$$

For every $\delta > 0$ let $B(\hat{x},\delta)$ be the open ball of radius δ centered at $\hat{x}$. Fix $r > 0$ such that $\hat{\sigma}(B(\hat{z},r)) \cap B(\hat{z},r) = \hat{\sigma}^{-1}(B(\hat{z},r)) \cap B(\hat{z},r) = \emptyset$, and consider a Hölder continuous function $\psi : \hat{\Sigma} \to \mathbb{R}$ such that

$$\psi(\hat{x}) = \begin{cases} 1 & \text{if} \quad \hat{x} = \hat{z} \\ 0 & \text{if} \quad \hat{x} \notin B(\hat{z},r). \end{cases}$$

Let R be a d-by-d complex matrix with small norm such that all the minors of $\mathrm{id}+R$ are non-zero. Such a matrix does exist: indeed, the set of matrices with non-vanishing minors is open set in the Zariski topology, and so it is dense in the norm topology. Define $\tilde{A}(\hat{x},t) = \hat{A}(\mathrm{id}+\psi(\hat{x})R)$. Then

$$H^{s,\tilde{A}}_{(\hat{x},h^u(t)),(\hat{p},h(t))} = \hat{A}(\hat{p},h(t))^{-1} H^{s,\hat{A}}_{\hat{\sigma}(\hat{x},h^u(t)),\hat{\sigma}(\hat{p},h(t))} \hat{A}(\hat{x},h^u(t))(\mathrm{id}+R).$$

Moreover, as the perturbation does not affect $\hat{A}$ outside $B(\hat{z},r)$,

$$H^{s,\tilde{A}}_{(\hat{x},h^u(t)),(\hat{p},h(t))} = A^{-1}\,\mathrm{id}\,A\,(\mathrm{id}+R) = (\mathrm{id}+R),$$

and so $H^{\tilde{A}}_t = (\mathrm{id}+R)$. As the Oseledets subspaces in ℓ coincide with the eigenspaces of A, the uniform twisting condition is equivalent to saying that the matrix of $H^{\tilde{A}}_t$ with respect to the eigenbase of A has all its minors different from zero. Thus $\tilde{A}$ is uniformly twisting and uniformly pinching. Making $R \to 0$, we can make the cocycle $\tilde{A}$ arbitrary close to $\hat{A}$. □

We need to prove that the set of simple cocycles with strong pinching and twisting is open.

Lemma 10.2. *The set of uniformly simple cocycles is open.*

Proof. First of all, existence of a dominated decomposition into 1-dimensional subspaces is equivalently to having, for each $l = 1,\dots,d$, stable and unstable cone fields, $C^s_l(t)$ and $C^u_l(t)$, such that $C^s_l(t)$ is an l-dimensional cone and $C^u_l(t)$ is a $d-l$ dimensional cone.

Let $E^s_l(t) = E^1_t + \cdots + E^l_t$ and $E^u_l(t) = E^l_t + \cdots + E^d_t$. If $\hat{A}$ is strongly pinching we can take the cones

$$C^s_{\epsilon,l}(t) = \{v = v^u_l + v^s_l, \quad \frac{\|v^u_l\|}{\|v^s_l\|} \le \epsilon \text{ where } v^u_l \in E^u_l(t) \text{ and } v^s_l \in E^s_l(t)\}$$

and

$$C^u_{\epsilon,l}(t) = \{v = v^u_l + v^s_l, \quad \frac{\|v^s_l\|}{\|v^u_l\|} \le \epsilon \text{ where } v^u_l \in E^u_l(t) \text{ and } v^s_l \in E^s_l(t)\}$$

and these are still, respectively, stable and unstable cone fields for every B close to $\hat{A}$. Moreover, if $\hat{B}$ is sufficiently close we can choose ϵ very small. Thus the spaces $E^i_{\hat{B}}$ varies continuously with $\hat{B}$.

By Remark 3.4 in [1], $\hat{A}_k \to \hat{A}$ in the space $H^{\hat{\alpha}}(\hat{M})$ implies that

$$H^u_{\hat{A}_k} \to H^u_{\hat{A}} \quad \text{and} \quad H^s_{\hat{A}_k} \to H^s_{\hat{A}}.$$

Then if $\hat{A}_k \to \hat{A}$ the spaces $E^i_{\hat{A}_k} \to E^i_{\hat{A}}$ and $H^s_{\hat{A}_k}{}^* H^u_{\hat{A}_k}{}^* \to H^s_{\hat{A}}{}^* H^u_{\hat{A}}{}^*$, so for k sufficiently large the transversality condition , strong twisting, is also verified. □

By Lemma 10.2, the subset of uniformly simple cocycles is open. By Theorem B every simple cocycle has simple Lyapunov spectrum. This concludes the proof of Theorem C.

Appendix A. Continuous maps are dense in $L^1(M, N)$

Let M be a normal topological space and N be a geodesically convex metric space (Definition 5.6). Denote by $\mathcal{F}$ the set of measurable maps $f : M \to N$. Given any regular σ-finite Borel measure μ on M, fix any point $\hat{0} \in N$ and define

$$L^1_\mu(M, N) = \{f \in \mathcal{F} : \int \operatorname{dist}_N \big(f(x), \hat{0}\big)\, d\mu(x) < \infty\}.$$

When μ is a finite measure, the choice of $\hat{0} \in N$ is irrelevant: different choices yield the same space $L^1_\mu(M, N)$.

The function $\operatorname{dist}_{L^1_\mu(M,N)} : L^1_\mu(M, N) \times L^1_\mu(M, N) \to \mathbb{R}$ defined by

$$\operatorname{dist}_{L^1_\mu(M,N)}(f, g) = \int d_N\big(f(x), g(x)\big) d\mu(x)$$

is a distance in $L^1_\mu(M, N)$. The special case $N = \mathbb{R}$ of the next proposition is well-known, but here we need the following more general statement:

Proposition A.1. *The subset of continuous maps $f : M \to N$ is dense in the space $L^1_\mu(M, N)$.*

We call $s : M \to N$ a *simple map* if there exist points $v_1, \dots, v_k \in N$ pairwise disjoint measurable sets $A_1, \dots, A_k \subset M$ with finite μ-measure such that

$$s(x) = \begin{cases} v_i & \text{if } x \in A_i \\ \hat{0} & \text{if } x \notin \cup_{i=1}^k A_i \end{cases}$$

Proposition A.1 is an immediate consequence of Lemmas A.2 and A.3 below.

Lemma A.2. *The set $\mathcal{S}$ of simple functions is dense in $L^1_\mu(M, N)$.*

Proof. Consider any $f \in L^1_\mu(M, N)$. Given $\epsilon > 0$, fix a set $K_0 \subset M$ with finite μ-measure and such that

$$\int_{M \setminus K_0} \operatorname{dist}_N(f(x), \hat{0})\, d\mu(x) \le \frac{\epsilon}{4}.$$

Let $\{v_1, \dots, v_i, \dots\}$ be a countable dense subset of N. The family

$$\big\{B\big(v_i, \frac{\epsilon}{\mu(K_0)}\big) : i \in \mathbb{N}\big\}$$

covers N and, consequently,

$$B_i = B\big(v_i, \frac{\epsilon}{2\mu(K_0)}\big) \setminus \bigcup_{j<i} B\big(v_i, \frac{\epsilon}{2\mu(K_0)}\big), \quad i \in \mathbb{N}$$

is a partition of N. Then $A_i = K_0 \cap f^{-1}(B_i)$, $i \in \mathbb{N}$ is a partition of K_0 into measurable sets. Fix $k \in \mathbb{N}$ large enough that

$$\int_{K_0 \setminus \bigcup_{i=1}^k A_i} \operatorname{dist}_N(f(x), \hat{0})\, d\mu(x) \le \frac{\epsilon}{4}.$$

Now define $s : M \to N$ by

$$s(x) = \begin{cases} v_i & \text{if } x \in A_i \text{ for } i = 1, \dots, k \\ \hat{0} & \text{if } x \notin \cup_{i=1}^k A_i. \end{cases}$$

Then

$$\int_{M \setminus \cup_{i=1}^k A_i} \operatorname{dist}_N(f(x), s(x))\, d\mu(x) = \int_{M \setminus \cup_{i=1}^k A_i} \operatorname{dist}_N(f(x), \hat{0})\, d\mu(x) \le \frac{\epsilon}{2}$$

and

$$\int_{\cup_{i=1}^k A_i} \operatorname{dist}_N(f(x), s(x))\, d\mu(x) \le \mu\big(\cup_{i=1}^k A_i\big) \frac{\epsilon}{\mu(K_0)} \le \frac{\epsilon}{2}.$$

Thus $\operatorname{dist}_{L^1_\mu(M,N)}(f, s) < \epsilon$, which proves the lemma. □

Lemma A.3. *For every $s \in \mathcal{S}$ and $\epsilon > 0$ there exists a continuous map $f : M \to N$ such that $\operatorname{dist}_{L^1_\mu(M,N)}(f, s) \le \epsilon$.*

Proof. Let A_i and v_i, $i = 1, \dots, k$ be as in the definition of the simple map s and $\tau \ge 1$ be as in Definition 5.6. Denote

$$L = \max\{d(v_i, \hat{0}) : i = 1, \dots, k\}.$$

For each $i = 1, \dots, k$, consider a compact set set $K_i \subset A_i$ such that $\mu(A_i \setminus K_i) < \epsilon/(4k\tau L)$. Since the K_i are are pairwise disjoint, and M is assumed to be normal, there exist pairwise disjoint open sets $B_i \supset K_i$, $i = 1, \dots, k$ with $\mu(B_i \setminus K_i) < \epsilon/(4k\tau L)$. In particular, we also have $\mu(A_i \setminus B_i) < \epsilon/(4k\tau L)$.

By the Urysohn lemma, there are continuous functions $\psi_i : M \to \mathbb{R}$, $i = 1, \dots, k$ such that

$$\psi_i(x) = \begin{cases} 1 & \text{if } x \in K_i \\ 0 & \text{if } x \notin B_i. \end{cases} \tag{53}$$

Now we use the assumption that N is geodesically convex. For each $i = 1, \dots, k$, fix $\lambda_i : [0, 1] \to N$ with $\lambda_i(1) = v_i$ and $\lambda_i(0) = \hat{0}$. Then define $f : M \to N$ by

$$f(x) = \begin{cases} \lambda_i(\psi_i(x)) & \text{if } x \in B_i \text{ with } i = 1, \dots, k \\ \hat{0} & \text{if } x \notin \bigcup_{i=1}^k B_i. \end{cases}$$

It is clear that f is continuous, because the B_i are open and pairwise disjoint. Moreover, $f(x) = s(x)$ if either

$$x \in \bigcup_{i=1}^k K_i \quad \text{or} \quad x \in \Big(\bigcup_{i=1}^k A_i \setminus \bigcup_{i=1}^k B_i\Big).$$

All the other values of x fall into some of the following cases:

(1) $x \in A_i \cap \big(B_j \setminus K_j\big)$ for some i and j: then

$$\begin{aligned} d_N(f(x), s(x)) &\le d_N(\lambda_j(\psi_j(x)), \hat{0}) + d_N(\hat{0}, v_i) \\ &\le \tau d_N(v_j, \hat{0}) + d_N(\hat{0}, v_i) \le 2\tau L. \end{aligned}$$

(2) $x \in A_i \setminus \cup_{j=1}^k B_j$ for some i: then

$$d_N(f(x), s(x)) = d(\hat{0}, v_i) \le L.$$

(3) $x \in B_j \setminus \cup_{i=1}^k A_i$ for some j: then

$$d_N(f(x), s(x)) = d_N(\lambda_j(\psi_j(x)), \hat{0}) \le \tau d_N(v_j, \hat{0}) \le \tau L.$$

In either case, x belongs to the set

$$\bigcup_{i=1}^{k}(A_i \setminus B_i) \cup \bigcup_{j=1}^{k}(B_j \setminus K_j)$$

which, by construction, has μ-measure bounded by $\epsilon/(2\tau L)$. So,

$$\mathrm{dist}_{L^1_\mu(M,N)}(f,s) \leq \frac{\epsilon}{2\tau L} \max\left\{d_N(f(x),s(x)) : x \in M\right\} \leq \epsilon,$$

as claimed. □

References

[1] A. Avila, J. Santamaria, and M. Viana. Cocycles over partially hyperbolic maps. Preprint www.preprint.impa.br 2008.

[2] A. Avila, J. Santamaria, and M. Viana. Holonomy invariance: rough regularity and applications to Lyapunov exponents. *Astérisque*, 358:13–74, 2013.

[3] A. Avila and M. Viana. Simplicity of Lyapunov spectra: a sufficient criterion. *Port. Math.*, 64:311–376, 2007.

[4] A. Avila and M. Viana. Extremal Lyapunov exponents: an invariance principle and applications. *Invent. Math.*, 181(1):115–189, 2010.

[5] L. Backes and A. Kocsard. Cohomology of dominated diffeomorphism-valued cocycles over hyperbolic systems. *Ergodic Theory Dynamical Systems*.

[6] C. Bonatti, X. Gómez-Mont, and M. Viana. Généricité d'exposants de Lyapunov non-nuls pour des produits déterministes de matrices. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 20:579–624, 2003.

[7] C. Bonatti and M. Viana. Lyapunov exponents with multiplicity 1 for deterministic products of matrices. *Ergod. Th. & Dynam. Sys*, 24:1295–1330, 2004.

[8] C. Bonatti and A. Wilkinson. Transitive partially hyperbolic diffeomorphisms on 3-manifolds. *Topology*, 44:475–508, 2005.

[9] M. Brin and Ya. Pesin. Partially hyperbolic dynamical systems. *Izv. Acad. Nauk. SSSR*, 1:177–212, 1974.

[10] C. Castaing and M. Valadier. *Convex analysis and measurable multifunctions*. Lecture Notes in Mathematics, Vol. 580. Springer-Verlag, 1977.

[11] H. Furstenberg. Non-commuting random products. *Trans. Amer. Math. Soc.*, 108:377–428, 1963.

[12] H. Furstenberg. Boundary theory and stochastic processes on homogeneous spaces. In *Harmonic analysis in homogeneous spaces*, volume XXVI of *Proc. Sympos. Pure Math. (Williamstown MA, 1972)*, pages 193–229. Amer. Math. Soc., 1973.

[13] H. Furstenberg and H. Kesten. Products of random matrices. *Ann. Math. Statist.*, 31:457–469, 1960.

[14] I. Ya. Gol'dsheid and G. A. Margulis. Lyapunov indices of a product of random matrices. *Uspekhi Mat. Nauk.*, 44:13–60, 1989.

[15] Y. Guivarc'h and A. Raugi. Products of random matrices : convergence theorems. *Contemp. Math.*, 50:31–54, 1986.

[16] M. Hirsch, C. Pugh, and M. Shub. *Invariant manifolds*, volume 583 of *Lect. Notes in Math.* Springer Verlag, 1977.

[17] V. I. Oseledets. A multiplicative ergodic theorem: Lyapunov characteristic numbers for dynamical systems. *Trans. Moscow Math. Soc.*, 19:197–231, 1968.

[18] O. Sarig. Symbolic dynamics for surface diffeomorphisms with positive entropy. *J. Amer. Math. Soc.*, 26:341–426, 2013.

[19] M. Viana. Almost all cocycles over any hyperbolic system have nonvanishing Lyapunov exponents. *Ann. of Math.*, 167:643–680, 2008.

[20] M. Viana. *Lectures on Lyapunov Exponents*. Cambridge University Press, 2014.

[21] M. Viana and K. Oliveira. *Foundations of Ergodic Theory*. Cambridge University Press, 2015.

IMPA – Estrada D. Castorina 110, Jardim Botânico, 22460-320 Rio de Janeiro, Brazil.
E-mail address: mpoletti@impa.br, viana@impa.br


xx
f2
f1
pp
zz
p
zk
z
x1
x2